

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
25 May 2001 (25.05.2001)

PCT

(10) International Publication Number  
**WO 01/36674 A2**

(51) International Patent Classification: C12Q 1/68,  
G01N 33/53, A61P 43/00

Dora [IE/GB]; 45 St. Anne's Avenue, Grappenhall, War-  
rington WA4 2PL (GB).

(21) International Application Number: PCT/GB00/04267

(74) Agent: GAINNEY, Laurence, David, Scott; Astrazeneca,  
Global Intellectual Property, P.O. Box 272, Mereside,  
Alderley Park, Macclesfield, Cheshire SK10 4GR (GB).

(22) International Filing Date:  
8 November 2000 (08.11.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
9926805.4 13 November 1999 (13.11.1999) GB

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(71) Applicant (*for all designated States except MG, US*): AS-  
TRAZENECA AB [SE/SE]; S-151 85 Sodertalje (SE).

(84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (*for MG only*): ASTRAZENECA UK LIM-  
ITED [GB/GB]; 15 Stanhope Gate, London W1Y 6LN  
(GB).

**Published:**

— Without international search report and to be republished  
upon receipt of that report.

(72) Inventors; and  
(75) Inventors/Applicants (*for US only*): BULL, John, Henry  
[GB/GB]; 25 Pownall Street, Macclesfield, Cheshire SK10  
1DF (GB). ELLISON, Gillian [GB/GB]; Alderley Park,  
Macclesfield, Cheshire SK10 4TG (GB). PASKINS, Lynn,

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: DIAGNOSTIC METHODS

(57) Abstract: The present invention derives from the identification of 26 gene transcripts (Markers) that exhibit aberrant expres-  
sion levels in prostate disorder tissues. The invention therefore relates to diagnostic techniques for the detection of human prostate  
disorders, such as cancer, by detecting one or more of these Markers, intermediates, precursors or products (mRNA, cDNA, genomic  
DNA, or protein). The invention is also directed to methods for identifying modulators of prostate disorders, which modulators, such  
as chemical compounds, antisense molecules and antibodies interact with and modulate any one of the Markers identified.

WO 01/36674 A2

PTS-0308

- 2 -

In early disease, where the cancer is confined to the prostate gland and has not spread beyond the gland, the tumour can potentially be removed by surgery. Surgery is a common treatment of prostate cancer. Transurethral resection is a procedure in which the cancer is cut from the prostate via the urethra. A radical prostatectomy is where the whole prostate and  
5 some of the tissue around the gland is removed. Radiation therapy is also used to treat early prostate cancer.

Once disease has become metastatic, its course is rapid and relatively predictable, leading to death within 2-5 years if untreated.

Prostate cancer commonly occurs in men above the age of 50 years. The incidence of  
10 prostate cancer rises sharply between the ages of 60 and 80 years, with more than 80% of all cases of the diseases being diagnosed in men over 65 and less than 1% in men under 50 years of age. The incidence and prevalence of prostate cancer have increased considerably throughout the world over the past two decades. The average age of detection is 72 in the UK and 66 in the US. Active screening programmes may lead to a lowering of the average age at  
15 first diagnosis.

Routine medical examinations for obstructive urinary symptoms may lead to an initial diagnosis of prostate cancer. When prostate cancer is asymptomatic, there are two tests that can be performed to detect the presence of cancer: digital rectal examination (DRE) and prostate specific antigen (PSA). PSA is an enzyme which is secreted almost exclusively by  
20 epithelial cells in the prostate; measurement of the quantity of PSA in the blood provides a more reliable indicator of the presence of prostate cancer than the DRE test. However, elevations in serum levels of PSA can also be found in patients with benign prostatic hyperplasia (BPH) and prostatitis. In some cases, where a patient has elevated levels of PSA in his blood, a third diagnostic technique is used called 'transrectal ultrasonography' (TRUS),  
25 which can further substantiate the presence of prostate cancer. In conjunction with taking a biopsy of the tissue, TRUS has been shown to increase the detection rate of prostate cancer two-fold, when compared to DRE alone. The combination of digital rectal examination, serum PSA level and transrectal ultrasound is currently the best available diagnostic tool (Gorgoulis VG. et al. Anticancer Res 19:2327-2348 (1999)).

30 It is common for the prostate gland to become enlarged with ageing, a condition called benign prostatic hyperplasia (BPH), or benign prostatic hypertrophy. Severe BPH can cause

- 4 -

nucleic acid targeted to the Marker mRNA; or more widely in the identification or development of chemical or hormonal therapeutic agents. The person skilled in the art is also capable of devising screening assays to identify compounds (chemical or biological) that modulate (activate or inhibit) one or other of the identified Marker, which compounds may  
5 prove useful as therapeutic agents in treating a prostate disorder. Monitoring the Markers could also be useful in identifying inhibitors, antagonists or biochemical signalling effects in high throughput screening.

According to a first aspect of the invention there is provided a method for diagnosing or prognosing or monitoring a prostate disorder comprising testing a biological sample for  
10 aberrant levels of one or more of the Markers selected from the group consisting of Marker 1 to Marker 26. Preferred Markers are numbers: 8, 13, 14, 15, 17, 18, 19, 20, 22, 24, 25 and 26.

The invention lies in the identification of these differentially expressed Markers in prostate disorders. Accordingly, the invention is directed to any diagnostic method capable of  
15 assessing the differential expression levels, relative to expression in control tissues, of one or more of the identified Markers, either alone or as a panel. In particular, such methods include assessment of mRNA transcript levels and/or protein levels. The presence of aberrant expression levels of one of the Markers indicating the presence of a prostate disorder.

In a preferred embodiment the diagnostic method involves testing for more than one of  
20 the Markers identified herein. As separate independent embodiments the diagnostic method may involve testing for 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or all of the Markers identified herein, optionally as part of a panel test of other gene Markers.

It will be apparent to the person skilled in the art that there are a large number of  
25 analytical procedures which may be used to detect the amount of any of the Marker products present in a test sample.

The test sample comprising nucleic acid or protein is conveniently any prostatic material or biological sample, including TURP chip, biopsy, excised prostate or part thereof, a sample of bone marrow aspirate, bone marrow biopsy, lymph node aspirate, lymph node  
30 biopsy, spleen tissue, fine needle aspirate, skin biopsy or organ tissue biopsy, particularly prostate gland tissue, invaded surgical margin, invaded lymph node, invaded lung, invaded

- 6 -

biomarkers for distinguishing cancer from BPH. Highly specific yet low abundance Markers could be invaluable if secreted in protein form, as they could permit non-invasive testing of body fluids.

According to a further aspect of the invention there is provided a method for  
5 distinguishing prostate cancer from BPH comprising testing a biological sample for aberrant levels of one or more of the Markers selected from the group consisting of Marker 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 17, 18, 20, 23, 25 and 26. Such a method might involve measuring the expression levels of one or more of these Markers in a biological sample taken from an individual, comparing the Marker expression level detected with control values or  
10 historical values and depending on the value detected determining whether or not the individual has prostate cancer or BPH.

Each of the Marker genes useful in the present invention are already in the public domain. However, the inventors are not aware of any prior art disclosing an association of any of the Marker genes with prostate cancer. Thus, although each of the Markers/genes  
15 identified herein are themselves already known, their association with prostate disorders leading to the present invention is unknown.

Marker 1 is the human smooth muscle protein, 22kDa. The gene coding for this protein has been cloned and sequenced (Thweatt et al., Biochem. Biophys. Res. Commun. 187:1-7 (1992)). These authors identify 22kD smooth muscle protein as a fibroblast or smooth  
20 muscle protein. There is no teaching in this paper of an association with prostate cancer. The sequence of 22kDa smooth muscle protein is disclosed in Thweatt et al. and is present in the EMBL database under accession number HS22SM. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HS22SM, unless stated otherwise or apparent from  
25 the context.

Marker 2 is the Rho GDP dissociation inhibitor protein. The gene coding for this protein has been cloned and sequenced (Leffers et al., Exp. Cell Res. 209:165-174 (1993)). These authors identify Rho GDP dissociation inhibitor protein as a keratinocyte protein. There is no teaching in this paper of an association with prostate cancer. The sequence of Rho  
30 GDP dissociation inhibitor protein is disclosed in Leffers et al., and is present in the EMBL database under accession number HSRHO1. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL

- 8 -

acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSD966, unless stated otherwise or apparent from the context.

Marker 7 is the human mRNA KIAA0120. This cDNA has been cloned and sequenced (Nagase et al., DNA Res. 2:37-43 (1995)). These authors identify KIAA0120 as a  
5 brain transcript. There is no teaching in this paper of an association with prostate cancer. The sequence KIAA0120 is disclosed in Nagase et al., and is present in the EMBL database under accession number HSORFF. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSORFF, unless stated otherwise or apparent from the context.

10 Marker 8 is the human ribosomal protein S25. The gene coding for this protein has been cloned and sequenced (Li et al., Gene 107:329-333 (1991)). These authors identify ribosomal protein S25 as a leukaemia cell line protein. There is no teaching in this paper of an association with prostate cancer. The sequence of human ribosomal protein S25 is disclosed in Li et al., and is present in the EMBL database under accession number HSRPS25.  
15 For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSRPS25, unless stated otherwise or apparent from the context.

Marker 9 is the human 80K-H protein (kinase C substrate). The gene coding for this protein has been cloned and sequenced (Sakai et al., Genomics 5:309-315 (1989)). These  
20 authors identify 80K-H protein (kinase C substrate) as a fibroblast and epidermal carcinoma cell protein. There is no teaching in this paper of an association with prostate cancer. The sequence of 80K-H protein (kinase C substrate) is disclosed in Sakai et al., and is present in the EMBL database under accession number HSG19P1A. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in  
25 the EMBL database under accession number HSG19P1A, unless stated otherwise or apparent from the context.

Marker 10 is the human alpha-2-macroglobulin protease inhibitor protein. The gene coding for this protein has been cloned and sequenced (Kan et al., Proc. Natl. Acad. Sci. U.S.A. 82:2282-2286 (1985)). These authors identify alpha-2-macroglobulin protease  
30 inhibitor protein as a serum protein. There is no teaching in this paper of an association with prostate cancer. The sequence of alpha-2-macroglobulin protease inhibitor protein is disclosed in Kan et al., and is present in the EMBL database under accession number

- 10 -

referred to herein, refers to that in the EMBL database under accession number HSLB2A26, unless stated otherwise or apparent from the context.

5 Marker 15 is the human PRSM1 protein. The gene for PRSM1 has been cloned and sequenced (Scott et al., Gene 174:135-143 (1996)). These authors identify PRSM1 as a putative metallopeptidase. There is no teaching in this paper of any association with prostate cancer. The sequence of PRSM1 is disclosed in Scott et al. and is present in the EMBL database under accession number HSU58048. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSU58048, unless stated otherwise or apparent from  
10 the context.

Marker 16 is the human ribosomal large subunit L12 protein. The gene for ribosomal large subunit L12 protein has been cloned and sequenced (Chu et al., Nucleic Acids Res. 21:749-749 (1993)). These authors identify human ribosomal large subunit L12 as a ribosomal protein. There is no teaching in this paper of any association with prostate cancer.  
15 The sequence of human ribosomal large subunit L12 is disclosed in Chu et al. and is present in the EMBL database under accession number HSL12A. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSL12A, unless stated otherwise or apparent from the context.

20 Marker 17 is the human 7SK45 RNA. The gene for 7SK45 RNA has been cloned and sequenced (Murphy et al., Nucleic Acids Res. 14:9243-9260 (1986)). These authors identify human 7SK45 as a small cytoplasmic RNA. There is no teaching in this paper of any association with prostate cancer. The sequence of 7SK45 is disclosed in Murphy et al. and is present in the EMBL database under accession number HS7K45. For the purpose of this  
25 application, the gene sequences and sequence positions referred to herein, refers to that in the EMBL database under accession numbers: HS7K45, unless stated otherwise or apparent from the context.

Marker 18 is the human KIAA0588 protein (human protocadherin gamma). This protein is abundantly expressed in brain cells. The gene for this protein has been cloned and  
30 sequenced (Nagase et al., DNA Research. 5:31-39 (1998)). There is no teaching in this paper of an association with prostate cancer. The sequence of the KIAA0588 protein is present in the EMBL database under accession number AB011160. For the purpose of this application,

- 12 -

application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSSRP14A, unless stated otherwise or apparent from the context.

Marker 23 is bone small proteoglycan I (biglycan) protein. The gene for this protein has been cloned and sequenced (Fisher et al., J. Biol. Chem. 264:4571-4576 (1989)). These authors identify bone small proteoglycan I (biglycan) as a protein derived from human bone-derived cells. There is no teaching in this paper of an association with prostate cancer. The sequence of bone small proteoglycan I (biglycan) protein is disclosed in Fisher et al. and is present in the EMBL database under accession number HSHPGI. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSHPGI, unless stated otherwise or apparent from the context.

Marker 24 is the human KIAA0045 gene. The gene for this protein has been cloned and sequenced (Nomura N et al., DNA Res 1:223-229 (1994)). These authors identify KIAA0045 as a protein from the human immature myeloid cell line KG-1. There is no teaching in this paper of an association with prostate cancer. The sequence of KIAA0045 is disclosed in the EMBL database under accession number HSKG1C (accession no. D28476). For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number HSKG1C, unless stated otherwise or apparent from the context.

Marker 25 is b-cell receptor associated protein, also known as REA (repressor of estrogen activity). The gene for this protein has been cloned and sequenced (Montano et al., Proc Natl Acad Sci U.S.A. 96:6947-6952 (1999)). These authors identify REA as a protein derived from human breast. There is no teaching in this paper of an association with prostate cancer. The sequence of b-cell receptor associated /REA protein, is disclosed in Montano et al. and is present in the EMBL database under accession number AF150962. For the purpose of this application, the gene and amino acid sequences and sequence positions referred to herein, refers to that in the EMBL database under accession number AF150962, unless stated otherwise or apparent from the context.

Marker 26 is the cystatin B protein. The gene for cystatin B has been cloned and sequenced (Pennacchio et al., Science 271:1731-1734 (1996)). These authors identify cystatin B protein as associated with progressive myoclonus epilepsy disease. There is no teaching in

- 14 -

markers such as PSA, prostatic acid phosphatase, prostate-specific membrane antigen or others (e.g. those reviewed in Gao et al., Prostate 31:264-281 (1977)); markers associated with cell proliferation, invasiveness, tumour angiogenesis, metastasis or any other aspect of carcinogenesis; or markers for other diseases and conditions. Levels of Marker mRNA in the test sample can be detected by any technique known in the art. These include Northern blot analysis, reverse transcriptase-PCR amplification (RT-PCR), microarray analysis and RNase protection.

In one embodiment, levels of Marker RNA in a sample can be measured in a Northern blot assay. Here, tissue RNA is fractionated by electrophoresis, fixed to a solid membrane support, such as nitrocellulose or nylon, and hybridised to a probe or probes capable of selectively hybridising with the Marker RNA to be detected. The actual levels may be quantitated by reference to one or more control housekeeping genes. Probes may be used singly or in combination. This may also provide information on the size of mRNA detected by the probe. Housekeeping genes are genes which are involved in the general metabolism or maintenance of the cell, and are considered to be expressed at a constant level irrespective of cell type, physiological state or stage in the cell cycle. Examples of suitable housekeeping genes are: beta actin, GAPDH, histone H3.3 or ribosomal protein L13 (Koehler et al., Quantitation of mRNA by Polymerase Chain Reaction. Springer-Verlag, Germany (1995)).

To gauge relative expression levels, a control sample can be run alongside the test sample or, the test result/value can be compared to Marker expression levels expected in a normal or control tissue. These control values can be generated from prior test experiments using normal or control tissues, to generate mean or normal range values for each Marker.

In another embodiment, the Marker nucleic acid in a tissue sample is amplified and quantitatively assayed. The polymerase chain reaction (PCR) procedure can be used to amplify specific nucleic acid sequences through a series of iterative steps including denaturation, annealing of oligonucleotide primers (designed according to the published Marker sequence to be detected), and extension of the primers with DNA polymerase (see, for example, Mullis, et al., U.S. patent No. 4,683,202; Loh et al., Science 243:217 (1988)). In reverse transcriptase-PCR (RT-PCR) this procedure is preceded by a reverse transcription step to allow a large amplification of the number of copies of mRNA (Koehler et al., *supra*).

Other known nucleic acid amplification procedures include transcription-based amplification systems (TAS) such as nucleic acid based sequence application (NASBA) and 3SR (Kwoh et



- 16 -

A variety of methods are currently available for making arrays of biological molecules. The 'dot or slot blot' approach, whereby an ordered array of DNA is vacuum blotted using a manifold, or hand blotted by capillary action, onto a porous membrane, such as nylon or nitrocellulose has been around for many years (Maniatis et al., Molecular Cloning-A Laboratory Manual, First Edition, Cold Spring Harbor, (1982)). Methods for preparing a plurality of oligonucleotide sequences and for attaching these to solid supports at high density are also known in the art. For example, US Patent No. 4,562,157 describes a method of using photo-activatable cross-linking groups to immobilise pre-synthesised ligands on surfaces. Fodor et al. (Nature 364:555-556 (1993)) and US Patent No. 5,143,854 describe the 'light-directed chemical synthesis' method for synthesising ligands, including oligonucleotides, directly onto a substrate surface at the desired location. US 5,700,637 also describes methods for *in situ* synthesis of oligonucleotides on solid support surfaces. In addition, such methods for preparing microarrays can easily be automated. International Publication No. WO 95/35505 discloses an automated capillary dispensing device and method for applying biological macromolecules to solid supports. International Publication No. WO 97/44134 also describes devices for delivery of small volumes of liquid (which may contain biological macromolecules) in a precise manner to produce micro-sized spots on a solid surface to generate a microarray. Similarly, International Publication No. WO 98/10858 also describes an apparatus for the automated synthesis of molecular arrays. Techniques exist for applying the oligonucleotides to the array at high density and for example, techniques exist for applying well in excess of  $10^3$  distinct polynucleotides per  $1 \text{ cm}^2$ .

Microarray technology makes it possible to simultaneously study the expression of many thousands of genes in a single experiment. Analysis of gene expression in human tissue (e.g. biopsy tissue) can assist in the diagnosis and prognosis of disease and the evaluation of risk for disease. A comparison of levels of expression of various genes from patients with defined pathological disease conditions with normal patients enables an expression profile, characteristic of disease, to be created. There are currently two main approaches to analyse gene expression using microarrays. In the first approach, cDNA fragments, often generated by PCR, for each of the genes under study are attached to an array. Typically, mRNA isolated from the test samples (i.e. induced or un-induced) is reverse transcribed into cDNA with incorporation of a fluorescent label. The cDNA is sheared and hybridised to the array. If a control test sample is to be run at the same time, mRNA from this sample can be reverse

- 18 -

In transcript profiling, several or many mRNAs are detected in the same procedure. One or more of these mRNAs may be diagnostic of cancer cells (i.e prostate cancer) in a tissue sample. In one embodiment, combinations of probes can be used to classify the cancer cells into clinically relevant types, according to the complex expression pattern of Markers measured on the array. Such classification may help to define which tumours are growing aggressively, or harbour latent signs of aggression, or are less aggressive or benign. The array provides a quantitative measure of Marker RNAs. This is done by comparison of Marker RNA signal with control signal. In a preferred embodiment hybridisation signals generated are measured by computer software analysis of images on phosphorimage screens exposed to radioactively labelled tissue RNA hybridised to a microarray of probes on a solid support such as a nylon membrane. In another, quantities are measured by densitometry measurements of radiation-sensitive film (e.g. X-ray film), or estimated by visual means. In another embodiment quantities are measured by use of fluorescently labelled probe, which may be a mixture of tumour and normal RNA differentially labelled with different fluorophores, allowing quantities of Marker mRNAs to be expressed as a ratio versus the normal level. The solid support in this type of experiment is generally a glass microscope slide, and detection is by fluorescence microscopy and computer imaging.

The detection of specific interactions may be performed by detecting the positions where the labelled target sequences are attached to the array. Radiolabelled probes can be detected using conventional autoradiography techniques. Use of scanning autoradiography with a digitised scanner and suitable software for analysing the results is preferred. Where the label is a fluorescent label, the apparatus described, e.g. in International Publication No. WO 90/15070, US Patent No. 5, 143,854 or US Patent No. 5,744,305 may be advantageously applied. Indeed, most array formats use fluorescent readouts to detect labelled capture:target duplex formation. Laser confocal fluorescence microscopy is another technique routinely in use (M.J.Kozal et al., Nature Medicine 2:753-759 (1996)). Mass spectrometry may also be used to detect oligonucleotides bound to a DNA array (Little et al, Analytical Chemistry 69: 4540-4546, (1997)). Whatever the reporter system used, sophisticated gadgetry and software may be required in order to interpret large numbers of readouts into meaningful data (such as described, for example, in US Patent No. 5,800,992 or International Publication No. WO 90/04652).

- 20 -

RNA can be extracted at a later date. Proprietary reagents are available which allow tissue or cells to be conveniently stored for several days at room temperature and up to several months at 4°C (e.g. RNAlater, Ambion Inc., TX ). Prior to extraction, methods such as grinding, blending or homogenisation are used to dissipate the tissue in a suitable extraction buffer.

- 5 Typical protocols then use solvent extraction and selective precipitation techniques. Example 1 describes such a method.

In another embodiment, tissue is directly analysed for the presence of Marker nucleic acid. This can be by *in situ* hybridisation, where sections of tissue may be interrogated with specific probes to determine which morphological cell type in the sample displays a marker nucleic acid, such as sequences corresponding to Markers 1-26. *In situ* hybridisation typically  
10 comprises 3 steps. Firstly tissue is fixed, and sections are prepared by standard treatments known to those in the art (Polack and McGee, *In situ* hybridisation: principles and practice. Oxford University Press, 1998). Secondly, Marker mRNA and amplified Marker DNA can be detected by hybridisation with e.g. a biotin-, digoxigenin- or radio-labelled Marker probe,  
15 typically for 2-16 hours at 42°C in a suitable hybridisation buffer. A typical buffer might contain 50% formamide, 5% dextran sulphate, 2xSSC and 10-20ng of probe per 7µl (Herrington and McGee, Diagnostic Molecular Pathology. IRL Press, Oxford, 1992). The probe can be made of DNA or RNA. Lastly, following stringency washes, the probes in hybridisation complexes are detected with chromogenic or fluorescent reagents, which can be  
20 visualised by microscopy, or by autoradiography in the case of radiolabelled probes. Signal amplification systems using e.g. tyramide can be used to increase sensitivity (Polack and McGee, *supra*). *In situ* PCR using oligonucleotide probes complementary to the nucleic acid of any of Markers 1 - 26 is therefore envisaged. *In situ* hybridisation can follow *in situ* PCR, giving greater specificity (Polack and McGee, *supra*). Techniques for quantitation of signal,  
25 and quantification of positive cells in a section are available to the pathologist using image analysis. *In situ* Marker visualisation permits localisation of signal in mixed-tissue specimens commonly found in tumours, and is compatible with many histological staining procedures. In one embodiment, several probes can be differentially labelled and hybridised simultaneously to the same section, and detected using appropriate reagents. In another, serial  
30 sections from the same sample can be analysed with a panel of probes. Quantitation may involve comparison with one or more control housekeeping genes as discussed above.

- 22 -

cDNA) then the probe or primer sequence can hybridise to the sense or antisense strand. If however the target is mRNA (single stranded sense strand) the primer/probe sequence will have to be the antisense complement.

5 An example of a suitable hybridisation solution when a nucleic acid is immobilised on a nylon membrane and the probe nucleic acid is greater than 500 bases or base pairs is: 6 x SSC (saline sodium citrate), 0.5% SDS (sodium dodecyl sulphate), 100µg/ml denatured, sonicated salmon sperm DNA. The hybridisation being performed at 68°C for at least 1 hour and the filters then washed at 68°C in 1 x SSC, or for higher stringency, 0.1 x SSC/0.1% SDS.

10 An example of a suitable hybridisation solution when a nucleic acid is immobilised on a nylon membrane and the probe is an oligonucleotide of between 12 and 50 bases is: 3M trimethylammonium chloride (TMACl), 0.01M sodium phosphate (pH 6.8), 1mM EDTA (pH 7.6), 0.5% SDS, 100µg/ml denatured, sonicated salmon sperm DNA and 0.1 dried skimmed milk. The optimal hybridisation temperature ( $T_m$ ) is usually chosen to be 5°C below the  $T_i$  of  
15 the hybrid chain.  $T_i$  is the irreversible melting temperature of the hybrid formed between the probe and its target. If there are any mismatches between the probe and the target, the  $T_m$  will be lower. As a general guide, the recommended hybridisation temperature for 17-mers in 3M TMACl is 48-50°C; for 19-mers, it is 55-57°C; and for 20-mers, it is 58-66°C.

A suitable hybridisation protocol is described in Example 5, however, operable  
20 variations to this method will be apparent to the person skilled in the art.

Where the cDNA molecules of Markers 1 - 26 encode proteins or parts of cellular proteins, these may themselves act as prostate disease Markers. To detect proteins in tissue, cells, body fluids or extracts of these sample types, specific antibody can be used. These antibodies can be prepared using the Marker protein/polypeptides.

25 Methods of making and detecting labelled antibodies are well known (Campbell; Monoclonal Antibody Technology, in: Laboratory Techniques in Biochemistry and Molecular Biology, Volume 13. Eds: Burdon R et al. Elsevier, Amsterdam (1984)). The term antibody includes both monoclonal antibodies, which are a substantially homogeneous population, and polyclonal antibodies which are heterogeneous populations. The term also includes inter alia,  
30 humanised and chimeric antibodies. Monoclonal antibodies to specific antigens may be obtained by methods known to those skilled in the art, such as from hybridoma cells, phage display libraries or other methods. Monoclonal antibodies may be inter alia, human, rat or

- 24 -

that the nucleic acid sequence encoding the polypeptide is transcribed into RNA in the host cell transformed or transfected by the expression vector construct. The coding sequence may or may not contain a signal peptide or leader sequence for secretion of the polypeptide out of the host cell. Expression and purification of the Marker polypeptides can be easily performed using methods well known in the art (for example as described in Sambrook et al. *supra*).

The Marker polypeptides so produced can then be used to inoculate animals, from which serum samples, containing the specific antibody against the introduced Marker protein/polypeptide, can later be obtained.

Rodent antibodies may be humanised using recombinant DNA technology according to techniques known in the art. Alternatively, chimeric antibodies, single chain antibodies, Fab fragments may also be developed against the polypeptides of the invention (Huse et al., Science 256:1275-1281 (1989)), using skills known in the art. Antibodies so produced have a number of uses which will be evident to the molecular biologist or immunologist skilled in the art. Such uses include, but are not limited to, monitoring enzyme expression, development of assays to measure enzyme activity and use as a therapeutic agent. Enzyme linked immunosorbant assays (ELISAs) are well known in the art and would be particularly suitable for detecting the Marker proteins or polypeptide fragments thereof in a test sample.

The expression system described in Example 3 can produce protein for use as an antigen for the generation of antibodies for use in an ELISA assay to detect Marker protein in body fluids or by immunohistochemistry (as described in Example 4) or other means. In addition, an antibody could be used individually or as part of a panel of antibodies, together with a control antibody which reacts to a common protein, on a dipstick or similar diagnostic device.

Levels of Marker gene expression can also be detected by screening for levels of polypeptide (Marker protein). For example, monoclonal antibodies immunoreactive with a Marker protein can be used to screen a test sample. Such immunological assays can be done in any convenient format known in the art. These include Western blots, immunohistochemical assays and ELISA assays. Functional assays can also be used, such as protein binding determinations.

In another preferred embodiment antibodies directed against a Marker protein or proteins can be used, to detect, prognose, diagnose and stage prostate cancer or its precursor lesions, or related prostate disorders. Various histological staining methods known in the art,

- 26 -

Thus, according to another aspect of the invention there is provided a method for treating a patient suffering from a prostate disorder comprising administering to said patient an effective amount of an anti-sense molecule capable of binding to the mRNA of a Marker gene selected from the group consisting of: Marker 1 - Marker 26, and inhibiting expression  
5 of the protein product of the Marker gene.

Complete inhibition of protein production is not essential, indeed may be detrimental. It is likely that inhibition to a state similar to that in normal tissues would be desired.

This aspect of antisense therapy is particularly applicable if the prostate disorder is a direct cause of over-expression of the Marker gene(s) in question, although it is equally  
10 applicable if said Marker gene(s) indirectly cause the prostate disorder. Having identified the particular Marker genes (1 - 26) over-expressed in prostate disorders, and with knowledge of the gene and mRNA sequence the person skilled in the art is able to design suitable antisense nucleic acid therapeutic molecules and administer them as required.

Antisense oligonucleotide molecules with therapeutic potential can be determined  
15 experimentally using well established techniques. To enable methods of down-regulating expression of a Marker gene of the present invention in mammalian cells, an example antisense expression construct can be readily constructed for instance using the pREP10 vector (Invitrogen Corporation). Transcripts are expected to inhibit translation of the gene in cells transfected with this type of construct. Antisense transcripts are effective for inhibiting  
20 translation of the native gene transcript, and capable of inducing the effects (e.g., regulation of tissue physiology) herein described. Oligonucleotides which are complementary to and hybridisable with any portion of Marker gene mRNA are contemplated for therapeutic use. U.S. Patent No. 5,639,595, "Identification of Novel Drugs and Reagents", issued Jun. 17, 1997, wherein methods of identifying oligonucleotide sequences that display *in vivo* activity  
25 are thoroughly described, is herein incorporated by reference. Expression vectors containing random oligonucleotide sequences derived from Marker polynucleotides are transformed into cells. The cells are then assayed for a phenotype resulting from the desired activity of the oligonucleotide. Once cells with the desired phenotype have been identified, the sequence of the oligonucleotide having the desired activity can be identified. Identification may be  
30 accomplished by recovering the vector or by polymerase chain reaction (PCR) amplification and sequencing the region containing the inserted nucleic acid material. Antisense molecules can be synthesised for antisense therapy. These antisense molecules may be DNA, stable

- 28 -

analogous to quantitative RT-PCR already described. Marker gene amplification could also be detected *in situ* in cells by probe hybridisation. Fluorescently-labelled probes (used in Fluorescent in situ Hybridisation or FISH) are ideal for this purpose as they allow subnuclear localisation via microscopy, and permit estimates of gene copy number by subnuclear spot intensity or number (Mark et al. Exp Mol Pathol 66:170-8 (1999)). Quantitative analysis of DNA for amplifications of Marker genes can also be carried out by Southern analysis, a method which is widely known to those skilled in the art (Sambrook et al., *supra*). DNA can be extracted from clinical material using established methods (Sambrook et al., *supra*). The methods of the invention can therefore also be directed to measuring genomic DNA levels of one or more of the identified Markers.

If a Marker mRNA encodes a secreted protein, that protein is likely to be present in body fluids. Proteins, especially secreted proteins, can be quantified in blood, serum, urine, semen and other fluids. Specific antibodies can often detect abundant proteins in ELISA tests on body fluid samples without enrichment. Prostate-specific antigen (PSA) falls into this category. PSA is an important Marker produced by prostatic epithelial cells and almost always expressed in prostate cancer, though not exclusively so. Clinically significant levels of proteins such as PSA are defined by appropriate studies, and protein levels are typically given as ng per mL of sample. Serum protein tests with greater specificity for cancer are needed. Detection of rare proteins may require that the protein is concentrated by e.g. precipitation. Thus, in a further embodiment of the invention diagnosis or prognosis or prostate disorder, or stage monitoring of the prostate disorder or therapeutic efficacy assessment is performed by testing for aberrant levels of one or more Marker proteins of the invention, which Marker protein is a secreted protein, in a bodily fluid.

The inventors predict that Markers 9, 19 and 24 (at least) are secreted proteins.

All the essential materials and reagents required for detecting or monitoring prostate disorder Markers in a test sample may be assembled together in a kit. Such a kit may comprise one or more diagnostic cDNA probes or oligonucleotide primers together with control probes/primers. The kit may contain probes immobilised on a microarray substrate such as a filter membrane or silicon-based substrate. The kit may also comprise samples of total RNA derived from tissues of various physiological states, such as normal, BPH, confined tumour and metastatic tumour, for example, to be used as controls. The kit may also comprise appropriate packaging and instructions for use in the methods of the invention.

- 30 -

Elmer 7700). Preferred oligonucleotide primers for detection of Marker mRNAs are selected from Markers 1-26.

In another embodiment the kit comprises one or more antibodies specific for one or other of the Markers identified herein for use in immunohistochemical analysis.

5 In another embodiment the kit is an ELISA kit comprising one or more antibodies specific for one or other of the Markers identified herein.

In another aspect of the invention, one or more of the 26 Markers can also be used in biochemical assays to identify agents which modulate the activity of the Marker proteins. The design and implementation of such assays will be evident to the biochemist of ordinary skill.

10 The protein, particularly if it is a biochemical enzyme, may be used to turn over a convenient substrate whilst incorporating or losing a labelled component to define a test system. Test compounds are introduced into the test system and measurements made to determine their effect on enzyme activity. Such assays are useful to identify inhibitors of the enzyme which may prove valuable as therapeutic agents.

15 The inventors believe that Marker 19 (Factor V) is an example of a suitable biochemical enzyme that can be used in a suitable biochemical assay to identify modulators.

In a further aspect of the invention, each of the Markers can be used to characterise cell cultures in a screen for therapeutic agents, such as a high throughput screen. Effects of test compounds may be assayed by changes in mRNA or protein of any of Markers 1-26. As  
20 described above, cells (i.e. mammalian, bacterial etc) can be engineered to express one of the Markers identified herein.

Thus, according to a further aspect of the invention there is provided a method of testing potential therapeutic agents for the ability to suppress a prostate disorder phenotype comprising contacting a test compound with a cell engineered to express one of the Markers  
25 identified herein; and determining whether said test compound suppressed expression of the Marker.

Thus, according to a further aspect of the invention there is provided a screening assay or method for identifying potential anti-prostate disorder therapeutic compounds comprising contacting an assay system capable of detecting the effect of a test compound against  
30 expression levels of one or more of the Markers selected from the group consisting of: Marker 1 - Marker 26, with a test compound and assessing the change in expression level of the particular Marker under study.



- 32 -

Further features of the invention include:

A method of treatment of a patient suffering from a prostate disorder, comprising administration to the patient of a compound capable of reducing the transcription or expression of any one of Markers 1 - 26.

- 5 A method of treatment of a patient suffering from a prostate disorder, comprising administration to the patient an antisense nucleic acid molecule targeted against the mRNA of any one of Markers 1 - 26.

Use of an antisense nucleic acid molecule or an antibody directed against any one of Markers 1 - 26, in the manufacture of a medicament for treating a prostate disorder.

- 10 Each aspect of the invention involves detection or use of one or more of Markers 1 - 26. A preferred sub-group of Markers are those selected from the group consisting of Markers: 8, 13, 14, 15, 17, 18, 19, 20, 22, 24, 25 and 26.

The invention will be further described by way of the following non-limiting examples and figures in which data illustrating over-expression of markers is included:

- 15 **Figure 1.** Filter microarray data showing over-expression of Markers 1 - 18 and 20-25 in prostate cancer relative to normal prostate. Overexpression value for Marker 26 (not shown) was found to be at the same level as Marker 13. Values given are mean expression level for 9 prostate cancer samples in those samples where over-expression of 1.5-fold or more was detected. Expression is given relative to normal prostate level (i.e. compared to the mean of 3  
20 normal prostate datasets. These 3 datasets comprised 2 different RNA mixtures, each with at least 10 normal sample components).

**Figure 2.** Filter microarray data showing over-expression of Markers 1-5, 7-12, 17, 18, 20 and 23 in prostate cancer relative to BPH. Values given are mean expression level for 9 prostate cancer samples in those samples where over-expression of 1.5-fold or more was  
25 detected. Expression is given relative to BPH level (i.e. compared to the mean of 13 BPH datasets).

#### Example 1: Identification and Evaluation of Markers of Prostate Disease by cDNA

##### Microarray Analysis

- 30 The microarray analysis protocol described in the following example was developed as a means to determine the relative abundances of mRNA species that are expressed in various tissues. Microarray analysis was used to identify differentially expressed RNA species

- 34 -

treated with DNase I (also Gibco BRL), again as recommended by the manufacturer. For 1<sup>st</sup> strand label synthesis, 20µg total RNA was reverse transcribed into cDNA in the presence of radiolabelled nucleotide (<sup>33</sup>P), using Superscript II enzyme (Gibco BRL) according to manufacturer's instructions. Labelled RNA was purified using GFX columns (Amersham Pharmacia Biotech, St. Albans, UK) according to manufacturers instructions, and added to 10 ml/filter of Church hybridisation buffer described below.

Prior to hybridisation, filters were wetted briefly in 2xSSC, then incubated in 5ml prewarmed Church hybridisation solution (0.5M sodium phosphate, 7% SDS, 1mM EDTA, pH 7.2) at 65°C for 2-6 hours. Probe was denatured at 100°C for 5 min, placed on ice for at least 5 min, then added to 5ml fresh Church hybridisation buffer in the presence of the filter, and mixed by swirling the bottle with the lid on for even probe distribution. Hybridisation was carried out at 65°C for 12-16 hours. Filters were rinsed briefly in pre-warmed Church wash solution (40mM sodium phosphate, 1% SDS, pH 7.2), then incubated twice in the same solution at 65°C for 20 min. Filters were drained briefly, wrapped in Saran wrap (Dow Chemical Company, USA) and exposed to phosphor screens for 3-6 days (screens and cassettes supplied by Molecular Dynamics, CA). The screens were then scanned using a Storm 830 phosphorimager (Molecular Dynamics). Array Vision software (Incyte Pharmaceuticals, CA, USA) was used to visualise the hybridisation images and generate quantitative numbers for each spot.

Typically, for comparison of data from different arrays, data is generated as a value relative to an internal standard for each array. Following export of ArrayVision data into Microsoft Excel format, spot measurements for data were normalised to housekeeping gene hybridisation signals known to be constant or relatively constant, using a simple Excel macro. A total of 38 spot values (from 16 different clones) representing 12 different housekeeping genes were used to generate an average housekeeping measurement. The gene and EMBL accession number of the housekeeping genes used are listed in Table 1. Background values were not subtracted, but to eliminate spurious low-level signals indistinguishable from noise, a local background value from 3 blank spots out of every 7x7 array was taken, and every value <2-fold this value was ignored.

To generate representative values for normal prostate, data from 3 array hybridisation experiments to normal prostate RNA was normalised to the mean housekeeping gene value, then the mean of these 3 values was taken as the control dataset.

- 36 -

histopathological analysis. Table 2 lists the 26 genes we have identified as being Markers for prostatic disease.

Control number	Gene	NCBI genInfo identifier	Spots on Array
1	Human ribosomal protein S14 gene	g337498	2
2	Human ribosomal protein S14 gene	g337498	2
3	Human acidic ribosomal phosphoprotein P0	g190231	2
4	Human alpha-tubulin mRNA	g340020	2
5	Human mRNA for TEF-5 protein	g1848081	2
6	Human transcriptional enhancer factor	g339440	2
7	Human transcription factor RTEF-1	g1561727	2
8	Human hypoxanthine phosphoribosyltransferase	g184349	2
9	Human hypoxanthine phosphoribosyltransferase	g184349	2
10	Human mRNA for transcription factor, TEF3	g1403337	2
11	Human acidic ribosomal phosphoprotein P1	g190233	2
12	Human mRNA for ribosomal protein L19	g36127	4
13	Human hH3.3B gene for histone H3.3	g761715	4
14	Human pancreatic phospholipase A-2 (PLA- 2)	g190008	2
15	Human mRNA for ribosomal protein L19	g36127	4
16	Human acidic ribosomal phosphoprotein P1	g190233	2

5 Table 1. Names and identities of control housekeeping genes used to normalise microarray data. This set was identified from a wider set, by comparison of variability of candidate housekeeping gene signals over 13 filter hybridisation experiments as a group showing minimal variability.

- 38 -

prostate cancer compared to BPH, or in the case of Marker 19, not detected at all in normal prostate. In addition, Markers 6, 13, 14 and 25 were not detected in BPH.

**Example 2: Quantitative RT-PCR test**

5 RT-PCR is a convenient method for assaying the relative abundance of mRNA species expressed in various cells, tissues and organs.

Reproducibly accurate amplification of an mRNA can be achieved by firstly synthesising a DNA template from the mRNA using a reverse transcriptase, then by PCR using a heat-stable DNA polymerase (e.g. Taq polymerase). An optimised amplification  
10 protocol is essential for quantitation of nucleic acids by PCR because small differences in efficiency of the reaction can greatly influence the reaction rates, with a subsequent effect on PCR yield. Optimisation of PCR is not usually problematic (refer to: Koehler *supra*, and references therein). Quantitation of RT-PCR products can be done while the reaction products are building up exponentially, preferably following each round of amplification  
15 (known as real-time PCR). For quantitation, analysis is carried out by reference to one or more housekeeping genes which are also amplified by RT-PCR, often in a different reaction tube. Quantitation of RT-PCR product may be undertaken, for example, by gel electrophoresis visual inspection or image analysis, HPLC (Koehler et al., *supra*) or by use of suitable detection methods such as described above.

20 In practise, specific pairs of oligonucleotide primers corresponding to Markers 1-26 can be used to trigger amplification of specific marker mRNAs. The Marker mRNA is quantitated relative to a standard housekeeping mRNA using a real-time RT-PCR assay. For the purpose of this example, the housekeeping gene is ribosomal protein L19, but any such gene could be used.

25 Firstly, RNA is isolated from a clinical or tissue sample as in example 1. Aliquots of RNA are used in a reverse transcription reaction using random hexamers or oligo-dT as primers, and for example Moloney Murine Leukaemia Virus (MuLV) or Avian Myeloblastosis Virus (AMV) reverse transcriptase. A typical 20ml reverse transcription reaction comprises the following: 10 mM each dNTP, 50 mM Tris-HCl, pH8.3, 50 mM KCl,  
30 50 mM dithiothreitol, 0.5 mM spermidine, 10ng oligo-dT primer, up to 1µg RNA, dissolved in DEPC-treated water. This mixture is incubated at 65°C for 15 min, then chilled on ice for 5 min. 5 units of AMV reverse transcriptase and 10 units of RNase inhibitor are added. The

- 40 -

**Example 3: Generation of antibody specific to Protein encoded by Markers 1-26**

Antibody specific to Markers 1-26 could be used to detect protein products of these Markers in tissue samples and body fluids. Ideally for diagnosis, easily accessible samples are required, involving minimally invasive techniques. Typically, these samples include  
5 blood, plasma, serum, saliva, urine and semen. These fluids are far removed from prostate cells and, in a normal individual, would not be expected to contain mRNA of prostatic origin. However, protein may be transported across cell membrane barriers into body fluids. Prostate-specific antigen is one such protein, and is easily assayed by a simple blood test using specific antibody against the protein. For new markers, e.g. Markers 1-26, it may be  
10 necessary to generate novel antibody. This example describes a method of generating protein in vitro, which can then be used to raise antibody.

One way to generate protein is to express the cloned gene in *E. coli*. This requires that the cDNA, e.g. Markers 1-26 is cloned into a vector capable of expressing the encoded protein in the bacterial host. Optionally, it may be desirable to incorporate a molecular tag into the  
15 protein so that it can be easily purified. One such tag in wide use is the 6xHis tag. A protein with this tag is easily isolated from a cell culture extract by affinity chromatography, and can be used in relatively pure form to inoculate an animal for antibody generation.

Having subcloned the cDNA or cDNA fragment into an appropriate vector, e.g. pHAT10, 11, 12 or 20 (Clontech, Palo Alto, CA) and confirmed the integrity of the insert such as sequence  
20 and orientation. The subclone is cultured on a suitable scale, cells are harvested by centrifugation and then lysed, and the protein extract incubated with TALON resin according to manufacturer's instructions (Clontech, Palo Alto, CA). The resin is washed and recombinant protein is eluted by adjusting the pH or imidazole concentration. If desired, the 6xHIS tag can be enzymatically removed using a specific protease (Clontech). To generate a  
25 specific antibody, the purified protein is injected into a host animal, usually rabbit, sheep or goat. After boost injections, the serum is periodically collected and tested for antibody.

Polyclonal antibody can be purified from the serum using standard techniques, or the animal's spleen can be harvested for the production of hybridomas. Techniques for the production of antibodies, both polyclonal and monoclonal, are well known to those skilled in the art (Catty  
30 D (ed.) Antibodies: A Practical Approach. Vol 1 (1988), Vol2 (1989)).

- 42 -

A hybridisation probe can be generated from a synthetic oligonucleotide or a dephosphorylated restriction fragment sequence by addition of a radioactive 5' phosphate group from [ $\gamma$ - $^{32}$ P]ATP by the action of T4 polynucleotide kinase. 20 pmoles of the oligonucleotide are added to a 20 $\mu$ l reaction containing 100mM Tris, pH7.5, 10mM MgCl<sub>2</sub>, 0.1mM spermidine, 20mM dithiothreitol (DTT), 7.55 $\mu$ M ATP, 55 $\mu$ Ci [ $\gamma$ - $^{32}$ P]ATP and 2.5u T4 polynucleotide kinase (Pharmacia Biotechnology Ltd, Uppsala, Sweden). The reaction is incubated for 30 minutes at 37°C and then for 10 minutes at 70°C prior to use in hybridisation. Methods for the generation of hybridisation probes from oligonucleotides or from DNA and RNA fragments are described in Chapters 11 and 10 respectively in Sambrook et al. (*ibid*). A number of proprietary kits are also available for these procedures.

#### Filter preparation

The sample DNA could be isolated and run on an agarose gel and Southern blotted onto a nitrocellulose or nylon filter using standard techniques.

#### Hybridisation conditions

Filters containing the nucleic acid are pre-hybridised in 100ml of a solution containing 6x SSC, 0.1%SDS and 0.25% dried skimmed milk (Marvel™) at 65°C for a minimum of 1 hour in a suitable enclosed vessel. A proprietary hybridisation apparatus such as model HB-1 (Techne Ltd) provides reproducible conditions for the experiment.

The pre-hybridisation solution is then replaced by 10ml of a probe solution containing 6xSSC, 0.1% SDS, 0.25% dried skimmed milk (e.g. Marvel™) and the oligonucleotide probe generated above. The filters are incubated in this solution for 5 minutes at 65°C before allowing the temperature to fall gradually to below 30°C. The probe solution is then discarded and the filters washed in 100ml 6xSSC, 0.1% SDS at room temperature for 5 minutes. Further washes (1-3) are then made in fresh batches of the same solution at 30°C and then, optionally, in 10°C increments up to 60°C for 5 minutes per wash.

After washing, the filters are dried and used to expose an X-ray film such as Hyperfilm™ MP (Amersham International) at -70°C in a light-tight film cassette using a fast tungstate intensifying screen to enhance the photographic image. The film is exposed for a suitable period (normally overnight) before developing to reveal the photographic image of the radio-active areas on the filters. Related nucleic acid sequences are identified by the presence of a photographic image compared to totally unrelated sequences which should not

- 44 -

then chilled quickly on ice-water, is placed on the tissue section and a coverslip is sealed on to the slide using rubber solution. A suitable probe is a digoxigenin-labelled 300-600 base pair cDNA clone or fragment thereof, corresponding to one of Markers 1-26, prepared by one of several methods known to the art. The slides are then incubated in a sealed chamber for 16 h at 37°C. The rubber solution is peeled away and coverslips are removed by immersion of the slides in 2x SSC, 0.1% SDS. Slides are washed 4 times for 5 min each at room temperature, and twice for 10 min at 65°C, in 2x SSC, 0.1% SDS, and rinsed briefly in 2x SSC. Hybridisation is detected as follows: Nonspecific binding is blocked by incubating the slides in blocking buffer (0.1M Tris-HCL pH 7.5, 0.1M NaCl, 2mM Mg Cl<sub>2</sub>, 3% bovine serum albumin) for 10 min. Slides are then flooded with anti-digoxigenin-conjugated alkaline phosphatase (1/500 dilution of stock; Boehringer, Mannheim, Germany) in blocking buffer, and incubated 2 h at room temperature, then washed in blocking buffer 3 times for 3 min each. Slides are placed in buffer 2 (0.1 M Tris-HCl pH 9.5, 0.1 M NaCl, 50 mM MgCl<sub>2</sub>) for 10 min, then immersed in substrate buffer (100mL buffer 2 containing 25 mg levamisole, plus 35 mg nitroblue tetrazolium chloride dissolved in 277 mL 70% dimethylformamide, plus 17 mg 5-bromo-4-indoyl-phosphate dissolved in 222 mL 100% dimethylformamide) for 10-30 min in the dark. The slides are then immersed in 20 mM Tris-HCl pH7.5, 5 mM EDTA for 5 min, rinsed with tap water for 5-10 min, mounted in aqueous mountant and examined under a microscope.

Example 7. Screening for therapeutic agents capable of inhibiting expression of any of Markers 1-26.



Markers 1-26 may be employed in a process for screening compounds which either inhibit, promote or modulate the expression of Markers 1-26. Examples of potential Marker 1-26 agonists are small molecules such as organic molecules or peptides, antibodies or oligonucleotides which bind to Markers 1-26 and inhibit expression or activity. One assay for therapeutic agents uses cultured cells, and measures transcript abundance using a microarray as described in Example 1. Typical prostate cell lines such as LNCaP, PC-3 and DU145 are available from the European Collection of Cell Cultures, Salisbury UK. Cell culture is a standard technique well known to those practised in the art. Briefly, in one suitable example, a seed vial of LNCaP cells is inoculated into 500mL culture flasks containing 20-50ml RPMI medium containing 10% fetal calf serum and 1% glutamate (all components Gibco, UK).

Claims:

1. A method for diagnosing or prognosing or monitoring a prostate disorder comprising testing a biological sample from an individual for aberrant levels of one or more of the  
5 Markers selected from the group consisting of: Marker 1 to Marker 26.
2. A method as claimed in claim 1, wherein the Marker(s) are selected from the group consisting of Markers: 8, 13, 14, 15, 17, 18, 19, 20, 22, 24, 25 and 26.
- 10 3. A method for distinguishing prostate cancer from BPH comprising testing a biological sample for aberrant levels of one or more of the Markers selected from the group consisting of Marker 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 17, 18, 20, 23, 25 and 26.
4. A method as claimed in any of claims 1-3 wherein mRNA transcript levels and/or  
15 protein levels of the Marker(s) is/are measured.
5. A method as claimed in claim 4, wherein mRNA transcript levels are measured using reverse-transcriptase polymerase chain reaction (RT-PCR).
- 20 6. A method for measuring Marker mRNA transcript levels as claimed in claim 4, which method involves the use of one or more oligonucleotide probes each capable of selectively hybridising to nucleic acid of a Marker of interest to determine the expression level of said Marker of interest.
- 25 7. A method for measuring Marker protein levels as claimed in claim 4, which method involves the use of one or more antibodies each capable of selectively binding to a Marker protein or protein fragment of interest to determine the expression level of said Marker of interest.
- 30 8. Use of an antibody selective for a Marker protein selected from the group consisting of: Marker 1 - Marker 26, in an assay to monitor therapeutic efficacy.



9. A diagnostic kit for diagnosing or prognosing or monitoring a prostate disorder comprising, one or more diagnostic probe(s) and/or one or more diagnostic primer(s) and/or one or more antibodies capable of selectively hybridising or binding to one or more of the Markers 1 -26.
- 5
10. A screening assay for identifying potential anti-prostate disorder therapeutic compounds comprising contacting an assay system capable of detecting the effect of a test compound on expression levels of one or more of the Markers selected from the group consisting of: Marker 1 - Marker 26, with a test compound and assessing the change in  
10 expression level of the particular Marker under study.
11. A method of testing potential therapeutic agents for the ability to suppress a prostate disorder phenotype comprising contacting a test compound with a cell engineered to express one of the Markers identified herein; and determining whether said test compound suppressed  
15 expression of the Marker.
12. A compound or agent identified by the screening assay according to claim 10 or the method according to claim 11.
- 20 13. A method of treatment of a patient suffering from a prostate disorder, comprising administration to the patient of a compound capable of reducing the transcription or expression of any one of Markers 1 - 26.

Figure 1Figure 2

- 2 -

cctccaccct ggtccgctc cctcgcttg ctcaaccgag tgcctccgac cccctcctc 840  
agccctcccc caccacacag cccagcctcc tcggtctcct gtctcgttgc tgcctctgcc 900  
tgtgtgtggt gggagagagg ccgcagccag gcctctgctg cccttctgtt gccccccagg 960  
ttctatctcc ccgtcacacc cgaggcctgg cttcaggagg gaggcggagca gccattctcc 1020  
5 aggccccgtg gttgccccg gacgtgtgcy tctgctgctc cggggtgag ctggggtgtg 1080  
ggatgcacgg cctcgtgggg gccgggcccgt cctccagccc cgtgctccc tggccagccc 1140  
ccttgtcgtt gtcgggtccc tctaaccatg atgecttaac atgtggagt taccgtgggg 1200  
cctcactagc cctactccc tgtgtctgca tgagcatgtg gcctccccgt cccttccccg 1260  
gtggcgaacc cagtgaacca gggacacgtg ggggtgtgctg gtgctgctcc ccagcccacc 1320  
10 aatgcctggc cagcctgccc ccttccctgg acagggctgt ggagatggct ccggcggctt 1380  
ggggaaagcg aaattgcaa cactcaagtc acctcagta catccaggag gctgggtatt 1440  
gtctgcctc tgccttttct gtctcagcgg cagtgccag agcccacacc ccccaagag 1500  
ccctcgatgg acaggcctga cccaccccac ctggggccag ccaggagccc cgcctggggc 1560  
atcagtattt attgcctccg tccgtgccgt cctggggcca ctgggtggcg cctcttcccc 1620  
15 cagcctctca gtgccaccac ccccgccagc cttccctgac ccagccagga caaacaaggg 1680  
accaagtga cacattgtg agagccgtct cctataggtc ccccgcccca tccccggtgt 1740  
tgggtgtgtg tctgccaggc tcaggcagag gcgcctgtcc ctgcttcttt tctgaccggg 1800  
aaataaatgc ccctgaagg 1819

20  
<210> 3  
<211> 996  
<212> DNA  
<213> Homo sapiens

25  
<400> 3  
agcttctctg atccctagta taaacacttc agtggtcccc tttcagttct actactttga 60  
ccgcgatgat gtggctttga agaactttgc caaataacttt cttcaccaat ctcatgagga 120  
gagggaacat gctgagaaac tgatgaagct gcagaaccaa cgagggtggc gaattcttct 180  
30 tcaggatata aaggtgaaca aaagatccta ggggtgtcat acttcatcat ctggcagtg 240  
tcgggtatca gaaatcactt aaactagcaa ttgcccttat aaagtgatga tacactgggc 300  
ttttgccttt tgtgcttttt taggcttacc atctaaacta aattaggcaa atagtaattg 360  
cccttttgcc aaaaacgtggg ggttagagat gatgggcttg ctgacttcta ggttagttgg 420  
tagagatgca ttaacctatt ctcatcaga aaccagactg tgatgactgg gagagcgggc 480  
35 tgaatgcaat ggagtgtgca ttacatttgg aaaaaaatgt gaatcagtca ctactggaac 540  
tgcacaaact ggccactgac aaaaatgacc ccatgtgag tattggaacc ccaggaaata 600  
aatggaggaa atcatttgce ttagggtatg ggaagctgc ccactaactg tcttgcceca 660  
ttgttttgca gttgtgtgac ttcattgaga cacattacct gaatgagcag gtgaaagcca 720  
tcaaagaatt ggggtgaccac gtgaccaact tgcgcaagat gggagcggcc gaatttggct 780  
40 tggcgaata tctctttgac aagcacaccc tgggagacag tgataatgaa agctaagcct 840  
cgggctaatt tccccatagc cgtgggggtg cttccctggt caccaaggca gtgcatgcat 900  
gttgggggtt cctttacctt ttctataagt tgtacaaaa catccactta agttctttga 960  
ttgtaccat tccctcaaat aaagaaattt ggtacc 996

45  
<210> 4  
<211> 2353  
<212> DNA  
<213> Homo sapiens

50  
<400> 4  
gggttgagca caaggaggag aacgaccaca aagtcttcta cgggggtgac ctgaaagtgg 60  
actgtgtggc caccggggtt cccaatcccc agatctcctg gaggctccca gacgggagtc 120  
tggtgaactc cttcatgcag tcggatgaca gcggtggacg caccaagcgc tatgtcgtct 180  
55 tcaacaatgg gacactctac tttaacgaag tggggtatgag ggagggaagg gactacacct 240  
gctttgtctg aaatcaggtc ggggaaggacg agatgagagt cagagtcaag gtggtgacag 300  
cgccccccac catccggaac aagacttgc tggcggttca ggtgccctat ggagacgtgg 360  
tcaacttagc ctgtgaggcc aaaggagaac ccatgcccac ggtgacttgg ttgtcccaaa 420  
ccaacaaggat gatccccacc tctctgaga agtatcagat ataccaagat ggcactctcc 480  
60 ttattcagaa agcccagcgt tctgacagcg gcaactacac ctgcttggtc aggaacagcg 540  
cgggagagga taggaagacg gtgtggattc acgtcaacgt ccagccaccc aagatcaacg 600  
gtaaccccaa ccccatcacc actgtgcggg agatagcagc cgggggcagt cggaaactga 660  
ttgactgcaa agctgaagcc atccccaccc cgagggtgtt atgggctttt cccgagggtg 720  
tggttctgcc agctccatac tatgaaacc ggatcactgt ccatggcaac ggttccctgg 780

&lt;211&gt; 5086

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

5 <400> 6  
gcctgcgcgc cgcagccgc ctcgggcggg aggcgggagg cgggaggccg ggcccaggcc 60  
ggggcagccc ctccaccgca cgcctcctggg ccggtgcccc ggtccgagtc gccttcgcc 120  
tgcccccccc gccaatcccc cgcgcgggca gccccagcca ggtcccgccg ccaggccggc 180  
tcccccccc gctccgcccc cggagccgca gccccgcccc ccatcgccgt cgcctatgtg 240  
10 tggctccgc agccggcgct ggggacgcgc gcggccgaga ctctggcctg cagtcggcgc 300  
cgccgcggcc aggtgggtgt tggactctag accatgtgcc taggtagaag ttttccctt 360  
ctccgcagct ctgctcccc agcaacgctc gccacaccct tgttttgaga tctctctaa 420  
ggagcggaga gtttaatagg caagaaggaa gggagaagac agaaggaga cgctccccg 480  
tacggagaca gagggaaggg gggctccaaa gccgaaagag gaggctcccta cctgccacg 540  
15 ataccagtca gcccttgcca gcatccagcc atgggggata tgaaaacccc agattttgat 600  
gaccttctgg ctgcctttga catccagac cccaccagcc ttgatgccaa ggaggccatc 660  
cagacaccca gtgaggagaa tgagagtccc ctcaaaccct caggcatatg tatggatgaa 720  
agtgtgtcct tgtctcactc aggtatcgcc cccgatgtgc cggccgtgag tgtcattgtc 780  
aagaacacca gccgcagga gtcatttgaa gcggagaaag accacattac tcccagtctc 840  
20 ctacacaatg gattccgggg ctccagatct cctccagatc cccacaactg tgggaaattt 900  
gattctactt ttatgaatgg agacagtgc aggagtctcc ctggcaaaact ggagcctccc 960  
aagtcagagc cattaccac cttcaaccag ttcagtccaa tctccagccc agaactgag 1020  
gatcccatca aagataaacg atttgggata aagcccaaac actctgacag ttatttccca 1080  
ccccctcttg ggtgcggggc tgtggggagg ccagtcctgg aggtcttggc taagtttccg 1140  
25 gttccagagc tgcataatgt tgatcatttt tgaagaaag aacccaagcc agaaccctcg 1200  
cccttgggga gccagcagga acacgagcaa agtgggcaga acacagtgga acctcacaag 1260  
gatccggatg ccaactcgatt cttcggggaa gctttggagt tcaacagcca tcttagcaac 1320  
agtattggag agtccaagg gcttgcggg gagcttggta cctgtctatc agtccccctt 1380  
aggcagcgtc taaagccagc tcattccaag ctgtcctctt gtgtggcagc ctgtgtggcc 1440  
30 ttgcaggcca aaagagtggc tagtgtcact aaggaggatc agcctggcca cacaaggat 1500  
ctctcagggc ccaactaaaga gatttctaaa ggtagcccca aaatgcccaa gtcaccaaag 1560  
agtccccgga gccctctgga ggccactaga aaaagtatca agccatcgga cagccctcgt 1620  
agcatctgca gtgacagcag cagcaaggc tcaccgtctg tggctgccag ctccccacca 1680  
gcaattccca aagtgagaat caaaaccatt aagacatcat caggggaaat caaacggact 1740  
35 gtcacaagga tcttgccaga tctgatgat ccaagtaagt cccctgttgg gtcacctcta 1800  
gggagcgcca ttgcagaggg cccagcgag atgccagggg atgaggtgcc tgtggaagag 1860  
cactttcctg aggcagggca aaattcaggg agccccaggg gggccaggaa aggggacgag 1920  
agcatgacaa aggcagtgta ctgctcact cccagctgca gttctgggcc cgggtccca 1980  
aagggggctg cccagggctc acagacaggc aagaagcaac agagcacagc actgcaggca 2040  
40 tccaccctgg cccctgccaa cctcctcccc aaagccgtgc acttggccaa cctgaacctc 2100  
gtccccaca gtgtgtgtgc atcagtgaac gccaaagtct cagtgcagg acggagccag 2160  
ccacagctta cacaatgtc ggtgcccctg gtccaccagg tgaaaaaggc tgcctcactg 2220  
attgtagagg tcttcaacaa ggtccttcac agctccaacc ccgtgccctt ctatgcgcca 2280  
aatctcagcc cgcctgcgga cagcaggatc cacgtgccgg ccagtgggta ctgctgcctg 2340  
45 gagtgtggag acgcaattgc cttagagaag agcctgagcc agcactatgg ccggcggagc 2400  
gtccacattg aggtactgtg cacactgtgc tccaagacgc tgctcttctt caacaagtgc 2460  
agcctgtctc ggcacgccc tgaccacaag agcaaggggc tcgtcatgca gtgttccag 2520  
ctgctggtga agcctatctc tgcggaccaaa atgttcgtgt cggccccctg gaactccacg 2580  
gcaccagcag cccagcccc ttcatcctct cccaaacatg gcctcacttc gggcagtgcc 2640  
50 agtccccctc ctccagcctt gccactctac ccagaccctg tgaggctcat ccggtactca 2700  
atcaagtgtc ttgaatgtca caagcagatg cgggactaca tggctcctggc tgcacatttc 2760  
cagaggacaa cagaggagac agaggggctg acctgcccag tatgccagat gctgtgcc 2820  
aaccagtga gtttctgtgc ccaccagcg attcatgcac acaagteccc ctactgtgc 2880  
ccggagtgtg gggctcctctg ccgctctgcc tacttccaga cccatgtaaa ggagaattgc 2940  
55 ctgcactatg cccgcaaggt gggctacagg tgcattccact gtggtgtcgt ccacctgacc 3000  
ttggccttgc tgaaaagcca catccaggag cgacactgcc aggttttcca caaatgtgca 3060  
ttctgcccc tggccttcaa gactgccagc accacagtgc cccccagcac 3120  
ccccccagc cccacagacc ctcccagctc atttataagt gctcctgtga aatggctctc 3180  
aacaagaaga ggcacattca gcagcatttt taccagaatg tcagcaagac gcaggtgggc 3240  
60 gtcttcaagt gccctgagtg cccactcttg ttcgtgcaga agccggagtt gatgcaaac 3300  
gtcaagagca cccacggtgt tcccgaagt gtggacgagc tgtcaaacct ccagtcttca 3360  
gcggacacat cctcaagccg cctggtctct cgagttccca ctgagccacc agccactagt 3420  
gtggctgtc ggagcagctc cctgccttct ggccgctggg gtaggcctga agccaccgc 3480  
aggggtggaag ccaggccgcg gctgaggaa actggctgga cctgccagga gtgccaggag 3540

- 6 -

<400> 6  
tttttttttt tttttttgtcc gacatcttga gacgaggtcg cggtgtctgc tgctattctc 60  
cgagcttcgc aatgccgcct aaggacgaca agaagaagaa ggacgctgga aagtcggcca 120  
agaaagacaa agacccagtg aacaaatccg ggggcaaggc caaaaagaag aagtgggtcca 180  
5 aaggcaaatg tcgggacaag ctcaataact tagtcttgtt tgacaaagct acctatgata 240  
aactctgtaa ggaagtcccc aactataaac ttataacccc agctgtggtc tctgagagac 300  
tgaagattcg aggtctccctg gccagggcag cccttcagga gctccttagt aaaggactta 360  
tcaaactggt ttcaaagcac agagctcaag taatttacac cagaaatacc aagggtggag 420  
atgctccagc tgctggtgaa gatgcatgaa taggtccaac cagctgtaca tttggaaaaa 480  
10 taaaacttta ttaaatc 497

<210> 9  
<211> 2056  
15 <212> DNA  
<213> Homo sapiens

<400> 9  
20 ggaaccgagg ctgctggaca agaggggtgc ggtggatact gacctttgct ccggcctcgt 60  
cgtgaagaca cagcgcatct ccccgctgta ggctttctcc acagaaccgg ttccgggcct 120  
cagagcgtct ggtgagatgc tgttgccgct gctgctgctg ctaccatgt gctgggccgt 180  
ggaggtcaag agggcccggg gcgtctccct caccaatcat cacttctacg atgagtccaa 240  
gcctttcacc tgcctggagc gttcggccac catccattt gatcaggtca acgatgacta 300  
25 ttgcgactgc aaagatggct ctgacgagcc aggcacggct gcctgtccta atggcagctt 360  
ccactgcacc aacactggct ataagccct gtatatcccc tccaaccggg tcaacgatgg 420  
tgtttgtgac tgctgcgatg gaacagacga gtacaacagc ggcgtcatct gtgagaacac 480  
ctgcaaagag aagggccgta aggagagaga gtccctgcag cagatggcgg aggtcaccgg 540  
cgaagggttc cgtctgaaga agatccctat tgaggactgg aagaaggcac gggaggagaa 600  
gcagaaaaag ctcatgtgac tacaggctgg gaagaagtct ctggaagacc aggtggagat 660  
30 gctgcggaca gtgaaggagg aagctgagaa gccagagaga gaggccaaag agcagaccca 720  
gaagctgtgg gaagagcagc tggctgctgc caaggcccaa caggagcagg agctggcggc 780  
tgatgccttc aaggagctgg atgatgacat ggacgggacg gtctcgggtg ctgagctgca 840  
gactcaccgg gagctggaca cagatgggga tggggcgttg tcagaagcgg aagctcaggg 900  
cctcctcagt ggggacacac agacagacgc cactcttttc tacgaccgag tctgggcccgc 960  
35 catcagggac aagtaccggt ccgaggcact gcccaccgac cttccagcac cttctgcccc 1020  
tgacttgacg gagcccaagg aggagcagcc gccagtcccc tcgtcgcccc cagaggagga 1080  
ggaggaggag gaggaggagg aagaagaggc tgaagaagag gaggaggagg aggattccga 1140  
ggaggcccca ccgccactgt caccgccgca gccggccagc cctgctgagg aagacaaaat 1200  
gccgccttac gacgagcaga cgcaggcctt catcgatgct gcccaggagg cccgcaacaa 1260  
40 gttcgaggag gccgagcggg cgctgaagga catggaggag tccatcagga acctggagca 1320  
agagattttt tttgactttg gccccaacgg ggagtttctg tacctgtaca gccagtgtca 1380  
cgagctcacc accaacgaat acgtctaccg cctctgcccc ttcaagcttg tctcgagaa 1440  
acccaaactc gggggctctc ccaccagcct tggcacctgg ggctcatgga ttggccccga 1500  
ccacgacaag ttcagtgcga tgaagtatga gcaaggcacg ggctgctggc agggcccaaa 1560  
45 ccgctccacc accgtgcgcc tctgtgcgg gaaagagacc atggtgacca gcaccacaga 1620  
gcccagtcgc tgcgagtacc tcatggagct gatgacgcca gccgctgcc cggagccacc 1680  
gcctgaagca cccaccgaag acgaccatga cgagctctag ctggatgggc gcagagaacc 1740  
tcaagaaggc atgaagccag cccctgcagt gccgtccacc cgccctctg ggctgctctg 1800  
tggctctgtt gccctcctct gtggcgccag gacctttgtg gggcttcgtg ccctgctctg 1860  
50 gggccccaggc ggggctggtc cacattccca ggccccaaac gcctccaaag atgggtaaag 1920  
gagcttgccc tccctgggccc cccacacttg gtgactcgcc ccaccacccc cagccctgtc 1980  
cctgccaccc ctccctagtgg ggactagtga atgacttgac ctgtgacctc aatacaataa 2040  
atgtgatccc ccaccc 2056

55 <210> 10  
<211> 4577  
<212> DNA  
60 <213> Homo sapiens

<400> 10  
gctacaatcc atctggtctc ctccagctcc ttctttctgc aacatgggga agaacaaact 60  
ccttcatecca agtctggttc ttctcctctt ggtcctcctg cccacagacg cctcagctctc 120  
tggaaccg cagtataagg ttctgggtccc ctccctgctc cactactgaga cactgagaa 180

- 8 -

5 ctccttgaaa tacaatatc tcccagaaaa ggaagagttc ccctttgctt taggagtgc 4080  
 gactctgcct caaacttgtg atgaacccaa agcccacacc agcttccaaa tctccctaag 4140  
 tgtcagttac acagggagcc gctctgcttc caacatggcg atcggtgatg tgaagatgg 4200  
 ctctggttc attccctcga agccaacagt gaaaatgctt gaaagatcta accatgtgag 4260  
 ccggacagaa gtcagcagca accatgtctt gattacctt gataaggtgt caaatcagac 4320  
 actgagcttg tctctcacgg ttctgcaaga tgtcccagta agagatctca aaccagccat 4380  
 agtgaaaagtc tatgattact acgagacgga tgagtttgca atcgctgagt acaatgctcc 4440  
 ttgcagcaaa gatcttgaa atgcttgaa accacaaggc tgaaaagtgc tttgctggag 4500  
 10 tctgttctc tgagctccac agaagacacg tgtttttgta tctttaaaga cttgatgaat 4560  
 aaacactttt tctggtc 4577

&lt;210&gt; 11

&lt;211&gt; 1651

15 &lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

20 ggagcccccg ccctgggatt cccaggtgtt ttcatttggg gatcagcact gaacacagaa 60  
 gagtcatgac ggagtttggg ctgagctggg ttttcttgt tgctattttt aaagggtgcc 120  
 agtgtgaggt gcagctgggt gagtctgggg gagacttggg ccagcctggg gggtccctga 180  
 gactctcttg tgcagcctct ggattcacct tcagtagtta tgctatgcac tgggtccgcc 240  
 aggtccagg gaaggagctg aaatatgttt caggtattag tagtaatggg cgtagaacat 300  
 attatgcaaa ttctgtgaag ggcagattca ccatctccag agacaattcc aagaacacgt 360  
 25 tgtatcttca aatgggcagc ctgagagctg aggacatggc tgtgtattac tgtgtgtccg 420  
 ggggaatcta tgatagtagt ggtccctttg actactggg ccagggaacc ctggtcaccg 480  
 tctcctcagc atccccgacc agccccagg tcttcccgct gagcctctgc agcaccacgc 540  
 cagatgggaa cgtggtcctc gctgctctg tccagggtt ctccccccag gagccactca 600  
 gtgtgacctg gagcgaaagc ggacagggcg tgaccgccag aaacttccca cccagccagg 660  
 30 atgctcccg ggacctgtac accacgagca gccagctgac cctgccggcc acacagtgcc 720  
 tagccggcaa gtccgtgaca tgccacgtga agcactacac gaatcccagc caggatgtga 780  
 ctgtgccctg cccagttccc tcaactccac ctaccccat tccctcaact ccacctacc 840  
 catctccctc atgtctccac ccccgactgt cactgcaccg accggccctc gaggacctgc 900  
 tcttaggttc agaagcgaac ctacgtgca cactgaccgg cctgagagat gcctcaggtg 960  
 35 tcaccttcac ctggacgccc tcaagtggga agagcgtgt tcaaggacca cctgagcgtg 1020  
 acctctgtgg ctgctacagc gtgtccagt tctgcccgg ctgtgccgag ccatggaacc 1080  
 atgggaagac ctctcactgc actgtgctt accccgagtc caagaccgg ctaaccgcca 1140  
 cctctcaaa atccggaaac acattccggc ccgaggtcca cctgtgccc ccgcccgcg 1200  
 aggagctggc cctgaacgag ctggtgacgc tgacgtgcct ggcacgcggc ttcagcccca 1260  
 40 aggacgtgct ggttcgctgg ctgcagggtt cacaggagct gccccgagc aagtacctga 1320  
 cttgggcac cggcaggag cccagccagg gcaccaccac ctctgctgtg accagcatal 1380  
 tgcgcgtggc agccgaggac tggaaagaag gggacacctt ctctgcatg gtgggccacg 1440  
 aggccctgcc gctggtcttc acacagaaga ccatcgaccg cttggcgggt aaaccaccc 1500  
 atgtcaatgt gtctgtgttc atggcggagg tggacggcac ctgctactga gccgcccgc 1560  
 45 tgtccccacc cctgaataaa ctccatgctc ccccaaaaaa aaaaaaaaaa aaaaaaaaaa 1620  
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 1651

&lt;210&gt; 12

50 &lt;211&gt; 1120

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 12

55 acttcttcgc accagggag cccacccac cagaacgcca agatgtccag caagcgggccc 60  
 aaagccaagg ccaccaagaa gcggccacag cgggccacat ccaatgtctt cgcaatgttt 120  
 gaccagtccc agatccagga gtttaaggag gctttcaaca tgattgacca gaaccgtgat 180  
 ggcttcattg acaaggagga cctgcacgac atgctggcct cgctggggaa gaaccccaca 240  
 gacgaatacc tggagggtcat gatgagcag gccccggggc catacaactt caccatgttc 300  
 60 ctccaccatgt ttggggagaa gctgaacggc acggaccccg aggatgtgat tcgcaacgcc 360  
 tttgctgct tgcagagga atcctcaggt ttcatccatg aggaccacct ccggaagctg 420  
 ctccaccacca tgggtgaccg ctccacagat gaggaagtgg acgagatgta ccgggaggca 480  
 cccgttgata agaaaggcaa cttaactac gtggagtcca cccgcctct caaacatggc 540  
 gccaaaggata aacacgacta ggccatcccc agccccctga caccagccc ccgccagtca 600

&lt;210&gt; 14

&lt;211&gt; 3612

&lt;212&gt; DNA

5 &lt;213&gt; Homo sapiens

&lt;400&gt; 14

10 aagcttaaaag tgaagacttg ctgaatttat ttcagcattt tactcctggc cattttccat 60  
aggaaggctc ctaatagtgt ctaactcctg aaagcagaaa gtgaccttca aaactgaccc 120  
atcttccctg gaatctctct aagagctgag ggattagggg tcttagagat ttgtgtatct 180  
ttgatggcac taaaatgaca tagagtttac atgtatacac ctggaaaagg tattaatagt 240  
tatcagatgg ccctcaaaag tctagaacat ggtagctgtg cattgaggaa acattctgca 300  
aaaaccattg tctcagaag aaacatccag ctctatatta attgtatgct cagacccatc 360  
cctgctgggt aggtcagcat ttgtacttgg ttgcatatt ataccattgt gtgacatcaa 420  
15 caatctgcaa gtacacattg ccgggctgcc tgtgcagaat ggcagctgct gggcctagag 480  
tctagtgcag gtacttctga agggatcatt ctactgaat atgtctgttc tcttctatgt 540  
acttctgac cctccagggg agctggatag agtggacctg aataagctaa acgagattga 600  
aggcacccta acaaaagcca aagatgaaat gaaggtcagc gatcttgata ggaaagtgtc 660  
tgacctggag aatgaagcca agaagcagga ggctgccatc atggactata accgagatat 720  
20 cgaggagatc atgaaggaca ttcgcaatct ggaggacatc aggaagacct taccatctgg 780  
ctgcttcaac acccctgcca ttgaaaagcc ctagtgtctt tagggctgga aggcagcatc 840  
cctctgacag gggggcagtt gtgaggccac agagtgcctt gacacaaaga ttacattttt 900  
cagaccccca ctctctgct gctgtccatc actgtccttt tgaaccagga aaagtcacag 960  
agtttaaaag gaagcaaat aaacatcctg aatcgggaac aaagggtttt atctaataaa 1020  
25 gtgtctcttc catcacgttg ctaccttacc cacacttccc tctgatttgc gtgaggacgt 1080  
ggcatcctac ttacgtacgt ggcataacac atcgtgtgag cccatgtatg ctggggtaga 1140  
gcaagttagc ctcccctgtc tcatcgatcc agcagaacct cctcagtctc agtactcttg 1200  
ttctataaag gaaaagtgtt gctactaaca gtacattgt gatggccagt atatccagtc 1260  
catggataaa gaaaatgcat ctgcatctcc tgcccctctt ccttctaagc aaaaggaaat 1320  
30 aaacatcctg tgccaaaggt attggtcatt tagaatgtcg gttagccatc atcagtgtct 1380  
ttagctatta tgagtgtagg aacttgagcc atccgtgggt caggatgcaa ttatttataa 1440  
aagtcccccag gtgaacatgg ctgaagattt ttctagtata ttaataattg actaggaaga 1500  
tgaacttttt ttcatgcttt tgggcagctg ataattttaa tctggatggg cagcttgac 1560  
tcaccaatag accaaaagac atcttttgat attcttataa atggaactta cacagaagaa 1620  
35 atagggatat gataaccact aaagttttgt tttaaaatc aaactaatc ttacagcttt 1680  
tttattagtt agtcttgaa ctagtgttaa gtatctggca gagaacagtt aatccctaag 1740  
gtcttgacaa aacagaagaa aaacaagcct cctcgtccta gtcttttcta gcaagggat 1800  
aaaacttaga tggcagcttg tactgtcaga atcccgtgta tccatttgtt cttctgttgg 1860  
agagatgaga catttgaccc tttagctcag ttttctctcg atgtttccat cttccagaat 1920  
40 ccctcaaaaa acattgtttg ccaaatcctg gtggcaata cttgcactca gtatttcaca 1980  
cagctgcaaa cgctatcgag ttctgcaact ttgtgattta aatccactct aaaccttccc 2040  
tctaagtcta gagggagac ccttacgtgg agtttcctag tgggcttctc aacttttgat 2100  
cctcagctct gtggttttaa gaccacagtg tgacagtccc ctgccacaca ccccttctct 2160  
cctaccaacc cacttttgag attcatatat agcctttaac actatgcaac ttgtactttt 2220  
45 gcgtagcagg ggctgggggt gggggaaaga aacctattat catggacaca ctggtgctat 2280  
taattatttc aaatttatat tttgtgtgta atgttttgtg ttttgtttat ccatgctata 2340  
gaacaaggaa tttatgtaga tatacttagt cctatttcta gaatgacact ctgttcaact 2400  
tgctcaattt ttctcttcca ctggcacaag tatctgaata cctccttccc tcccttctag 2460  
agttctttgg attgtactcc aaagaattgt gccttgtgtt tgcagcatct ccattctcta 2520  
50 aattaatata attgtcttcc tccacacca gccacgtaaa gaggttaact gggctctctt 2580  
ccattgcagt cctgatgac ctaacctgca gcacgggtgt tttacaatgt tccagagcag 2640  
gaacgccagg ttgacaagct atggtaggat taggaaagtt tgctgaagag gatctttgac 2700  
gccacagtgg gactagccag gaatgaggga gaaatgccct ttttggcaat tgttggagct 2760  
ggataggtaa gttttataag ggagtacatt ttgactgagc acttagggca tcaggaacag 2820  
55 tgctacttac tgggtggtag actgggagag gtggtgtaac ttagtctctg atgatccac 2880  
ttctgttttc catctgcttg ggatatacca gagtttacca caagtgtttt gacgatatac 2940  
tcttgagctt tcaactctgt ggcttctccc aggcctcttc tactatggca ggagatgtgg 3000  
tgtgtgttgg caaagttttc acgtcatcgt ttcttggtta gttcatttca ttaagtggct 3060  
acatcctaac atatgcattg gtcaagggtg cagcaaggag actgaagatt gactgccaag 3120  
60 ctagtgttggg tgaagttcac tccagcaagt ctgagccac aatggggtgg tttgggttgg 3180  
tttcttttta actttctttt tgttatattg ttttctctc cacctgtgtg gtatattttt 3240  
taagcagaat tttatttttt aaaataaaaag gttctttaca agatgatacc ttaattacac 3300  
tcccgaaca cagccattat tttattgtct agctccagtt atctgtattt tatgtaattg 3360  
aatgacagg atggctgctg cagaatgctg gttgacacag ggattattat actgctattt 3420

ccggaggtga agtcgggtgcc acttctgccc tggcccccga gatcggcccc ctgggtctgt 180  
 ctccaaaaaa agttgggtgat gacattgcca aggcaacggg tgactggaag ggcttgagga 240  
 ttacagtga actgaccatt cagaacagac aggccagat tgaggtggtg ccttctgcct 300  
 ctgccctgat catcaaagcc ctcaaggaa caccaagaga cagaaagaaa cagaaaaaca 360  
 5 ttaaacacag tgggaatatc acttttgatg agattgtcaa cattgctcga cagatgcggc 420  
 accgatcctt agccagagaa ctctctggaa ccattaaaga gatcctgggg actgccagtg 480  
 cagtgggctg taatgttgat ggccgccatc ctcatgacat catcgatgac atcaacagtg 540  
 gtgctgtgga atgcccagcc agttaagcac aaaggaaaac atttcaataa aggatcattt 600  
 gacaactggg ga 612  
 10  
 <210> 17  
 <211> 1100  
 <212> DNA  
 15 <213> Homo sapiens  
 <400> 17  
 ctgcagtatt tagcatgccc caccatctg caaggcattc tggatagtgt caaacagcc 60  
 ggaaatcaag tccgtttatc tcaaaacttta gcattttggg aataaatgat atttgctatg 120  
 20 ctggttaaat tagattttag ttaaatcttc tgctgaagct ctagtacgat aagcaacttg 180  
 acctaatgtt aaagttaga tttccttcag gtttatatag ctgtgcgcc gcctgggtac 240  
 ctccgagtgt agggcgatct ggctgcgaca tctgtcacc cttgatcgc cagggttgat 300  
 tggctgatc tggctggcta ggctgggtgt cccttctcc ctaccgctc catgtgctc 360  
 cctccgaag ctgcgcgctc ggtcgaagag gacgaccatc cccgatagag gaggaccgg 420  
 25 ctccggtcaa gggatataca gtatgctgc tccctgtc gaacctcaa acaagctctc 480  
 aaggtccatt tgtaggagaa cgtagggtag tcaagcttcc aagactccag acacatcaa 540  
 atgaggcgct gcatgtggca gtctgccttt ctttgaccc attaccatc taaggaccgg 600  
 tcttcgggtca aggggtatac agtagctgct ctcctctgct agaacctcca acaagctct 660  
 caaggtccat ttgtaggaga acgtagggtg gtcaagcttc caagactcca gacacatcca 720  
 30 aatgaggcgc tgcattgtggc agtctgcctt tcttttgacc cattaccat ctaagttaga 780  
 tgctttttta aatgtttttt aatttttaaa tttttaattt ttttcattat ttatttttta 840  
 tttttgagac ggaatctcggc tcactgtaac ctccacctcc cgggttcaag cgattctcct 900  
 gactcagcct cccgagtagc tgggattaca ggcgcgcgcc accatgccc gctaattttt 960  
 gtatttttag taaagacggg ttttcgcat gttagcagg gtggtcttga actcctgacg 1020  
 35 tcaggtgatc ctctgcctc gtgctgggat tgcaggcgtg agccaccgct cccggccctt 1080  
 agatgctttt taatcagcaa 1100  
 <210> 18  
 40 <211> 4747  
 <212> DNA  
 <213> Homo sapiens  
 <400> 18  
 45 taggaatgaa ccatccctgg gagtatgtgg aaacaacgga ggagctctga cttcccaact 60  
 gtcccatctt atgggcgaag gaactgctcc tgacttcagt gggttaagggc agaattgaaa 120  
 ataattctgg aggaagataa gaatgattcc tgcgcgactg caccgggact acaaagggtc 180  
 tgcctgctg ggaatcctcc tggggactct gtgggagacc ggatgcaccc agatacgcta 240  
 ttcagttccg gaagagctgg agaaaggctc taggggtggg gacatctcca gggacctggg 300  
 50 gctggagccc cgggagctcg cggagcgctg agtccgcac atcccagag gtaggacgca 360  
 gcttttcgcc ctgaatccgc gcagcgccag cttggtcacg gcgggcagga tagaccggga 420  
 ggagctctgt atggggggcca tcaagtgtca attaaatcta gacattctga tggaggataa 480  
 agtgaaaata tatggagtag aagtagaagt aaggacatt aacgacaatg cgccttactt 540  
 tctgtaaagt gaattagaaa taaaaattag tgaatgca gccactgaga tgcggttccc 600  
 55 tctacccac gcctgggatc cggatattcg gaagaactct ctgcagagct acgagctcag 660  
 cccgaacact cacttctccc tcatcgtgca aaatggagcc gacggtagta agtaccgca 720  
 attggtgctg aaacgcgccc tggaccgca agaaaaggct gctcaccacc tggctcttac 780  
 ggctccgac gggggcgacc cggtcgcac aggcaccgag cgcatccgag tgatggttct 840  
 ggatgcgaac gacaacgcac cagcgtttgc tcagcccag taccgcgca gcgttccgga 900  
 60 gaatctggcc ttgggcacgc agctgcttgt agtcaacgct accgaccctg acgaaggagt 960  
 caatgcggaa gtgaggtatt ccttcgggta tgtggacgac aaggcgggcc aagttttcaa 1020  
 actagattgt aattcaggga caatatcaac aataggggag ttggaccag aggagtcagg 1080  
 attctaccag atggaagtgc aagcaatgga taatgcagga tattctgcgc gagccaaagt 1140  
 cctgatcact gttctggacg tgaacgacaa tgcccagaa gtggtcctca cctctctcgc 1200



&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 19

5 gaattccgca gcccgagtg tggtagcag ctcggcaagc gctgcccagg tcctggggtg 60  
gtggcagcca gcgggagcag gaaaggaagc atgttcccag gctgcccag cctctgggtc 120  
ctggtggtct tgggcaccag ctgggtagcc tgggggagcc aaggagcaga agcggcacag 180  
ctaaggcagt tctacgtggc tgctcagggc atcagttgga gctaccgacc tgagcccaca 240  
aactcaagtt tgaatctttc tgtaactttc ttttaagaaa ttgtctacag agagtatgaa 300  
10 ccatatttta agaaagaaaa accacaatct accatttcag gacttcttgg gcctacttta 360  
tatgtgtaag tcggagacat cataaaagt cactttaaaa ataaggcaga taagcccttg 420  
agcatccatc ctcaaggaa taggtacagt aaattatcag aaggtgcttc ttaccttgac 480  
cacacattcc ctgaggagaa gatggacgac gctgtggctc caggccgaga atacacctat 540  
gaatggagta tcagtgaaga cagtggaccc accatgatg accctccatg cctcacacac 600  
15 atctattact cccatgaaaa tctgatcgag gatttcaact cggggctgat tgggcccctg 660  
cttatctgta aaaaaggagc cctaactgag ggtgggacac agaagacgtt tgacaagcaa 720  
atcgtgctac tatttgctgt gtttgatgaa agcaagagct ggagccagtc atcatcccta 780  
atgtacacag tcaatggata tgtgaatggg acaatgccag atataacagt ttgtgcccat 840  
gaccacatca gctggcatct gctgggaatg agctcggggc cagaattatt cctcattcat 900  
20 ttcaacggcc aggtcctgga gcagaacat cataaggtct cagccatcac cctgtcagt 960  
gtacatccca ctaccgcaaa tatgactgtg ggcccagagg gaaagtggat catatcttct 1020  
ctcaccctcaa aacattttga agctgggatg caggcttaca ttgacattaa aaactgccca 1080  
aagaaaacca ggaatcttaa gaaaataact cgtgagcaga ggcggcacat gaagaggtgg 1140  
gaatacttca ttgctgcaga ggaagtcat tgggactatg cactgtaat accagcgaat 1200  
25 atggacaaaa aatacaggtc tcagcatttg gataatttct caaaccat tggaaaacat 1260  
tataagaaag ttatgtacac acagtacgaa gatgagtcct tcaccaaaca tacagtgaat 1320  
cccaatatga aagaagatgg gattttgggt cctattatca gagcccaggt cagagacaca 1380  
ctcaaaatcg ttttcaaaaa tatggccagc cgtccctata gcattttacc tcatggagt 1440  
accttctcgc cttatgaaga tgaagtcaac tcttctttca cctcaggcag gaacaacacc 1500  
30 atgatcagag cagttcaacc aggggaaacc tatacttata agtggaaacat cttagagttt 1560  
gatgaaccca cagaaaatga tggccagtgc ttaacaagac catactacag tgacgtggac 1620  
atcatgagag acatcgccctc tgggctaata ggactacttc taatctgtaa gacgagatcc 1680  
ctggacaggg gaggaatata gagggcagca gacatcgaaac agcaggctgt gtttgctgtg 1740  
tttgatgaga acaaaaagctg gtacccttgag gacaacatca acaagttttg tgaaaatcct 1800  
35 gatgaggtga aacgtgatga ccccaagttt tatgaatcaa acatcatgag cactatcaat 1860  
ggctatgtgc ctgagagcat aactactctt ggattctgct ttgatgacac tgtccagtgg 1920  
cacttctgta tgttggggac ccagaatgaa attttgacca tccacttcac tgggactcca 1980  
ttcatctatg gaaagaggca tgaggacacc ttgaccctct tccccatgcy tggagaatct 2040  
gtgacggtca caatggataa tgttggaaact tggatgttaa cttccatgaa ttctagtcca 2100  
40 agaagcaaaa agctgaggct gaaattcagg gatgttaaat gtatcccaga tgatgatgaa 2160  
gactcatatg agatttttga acctccagaa tctacagtea tggctacag gaaaatgcat 2220  
gatcgtttag aacctgaaga tgaagagagt gatgctgact atgattacca gaacagactg 2280  
gtcgagcat taggaattag gtcattccga aactcatcat tgaaccagga agaagaagag 2340  
ttcaatctta ctgccctagc tctggagaat ggcactgaat tctgttcttc gaacacagat 2400  
45 ataattgttg gttcaaatla ttcttcccca agtaatatla gtaagttcac tgtcaataac 2460  
cttgacagaa ctacagaaagc ccttctctac caacaagcca ccacagctgg tccccactg 2520  
agacacctca ttggcaagaa ctcagttctc aattcttcca cagcagagca ttccagccca 2580  
tattctgaag acctataga ggatcctcta cagccagatg tcacagggat acgtctactt 2640  
tcacttggtg ctggagaatt cagaagtcaa gaacatgcta agcgtaaagg acctaaaggt 2700  
50 gaaagagatc aagcagcaaa gcacagggtc tcctggatga aattactagc acataaagtt 2760  
gggagacacc taagccaaga cactggttct ccttcgggaa tgaggccctg ggaggacctt 2820  
cctagccaag aactggttc tcttccaga atgaggccct gggaggaccc tctagtgat 2880  
ctgttactct taaaacaaag taactcatct aagattttgg ttgggagatg gcatttggct 2940  
tctgagaaag gtatgctatga aataatccaa gatactgatg aagacacagc tgttaacaat 3000  
55 tggctgatca gccccagaa tgctcacgt gcttggggag aaagcaccct tcttgccaac 3060  
aagcctggaa agcagagtgg ccacccaag tttcctagag ttagacataa atctctacaa 3120  
gtaagacagg atggaggaaa gagttagctg aagaaaagcc agtttctcat taagacacga 3180  
aaaaagaaaa aagacaaagc cacacacat gctcctttat ctccaggagc ctttaccct 3240  
ctaagaagtg aagcctacaa cacattttca gaaagaagac ttaagcatc gttgtgctt 3300  
60 cataaatcca atgaacatc tcttccaca gacctcaatc agacattgcc ctctatggat 3360  
tttggtgga tagctcact tctgacct aatcagaatt cctcaaatga cactggtcag 3420  
gcaagctgtc ctccaggtct ttatcagaca gtgccccag aggaacacta tcaaacatc 3480  
ccattcaag acctgatca aatgcactct acttcagacc ccagtcacag atcctcttct 3540  
ccagagctca gtgaaatgct tgagtatgac cgaagtcaca agtccttccc cacagatata 3600

ccgcccgtt ctactccgcc gcgggggtcg cagcggctgc cgcgccgtcc tcgagtttcc 60  
agcgtgagga ggaggctgag ggaggagagg cgcacgtgt tcgaggcgga gaccgagggg 120  
gagccccgcg cgcggcgctc ctcatgtcta tggacagtgc tatcaccctg tggcagttcc 180  
ttcttcagct cctgcagaag cctcagaaca agcacatgat ctgttggacc tctaagtatg 240  
5 ggcagtttaa gcttttgcag gcagaagagg tggctcgtct ctgggggatt cgcaagaaca 300  
agcctaacat gaattatgac aaactcagcc gagccctcag atactattat gtaagaata 360  
tcatcaaaaa agtgaatggt cagaagtgtg tgtacaagtt tgtctcttat ccagagattt 420  
tgaacatgga tccaatgaca gtgggcagga ttgagggtga ctgtgaaagt ttaaacttca 480  
gtgaagtcat cagcagttcc aaagatgtgg agaattggagg gaaagataaa ccacctcagc 540  
10 ctgggtccaa gacctctagc cgcaatgact acatacactc tggcttatat tcttcattta 600  
ctctcaactc tttgaactcc tccaatgtaa agcttttcaa attgataaag actgagaatc 660  
cagccgagaa actggcagag aaaaaatctc ctccaggagcc cacaccatct gtcatacaat 720  
ttgtcacgac accttccaaa aagccaccag ttgaacctgt tgctgccacc atttcaattg 780  
gccaagtat ttctccatct tcagaagaaa ctatccaagc tttggagaca ttggtttccc 840  
15 caaaactgcc ttccctggaa gcccacact ctgcctctaa cgtaatgact gcttttgcca 900  
ccacaccacc catttcgtcc ataccctctt tgcagggaacc tcccagaaca ccttcaccac 960  
cactgagttc tcacccagac atcgacacag acattgattc agtggcttct cagccaatgg 1020  
aacttccaga gaatttgtct ctggagccta aagaccagga ttcagtcttg ctagaaaagg 1080  
acaaagttaa taattcatca agatccaaag aacccaaagg gttaggactg gcaccacccc 1140  
20 ttgtgatcac gagcagtgat ccaagcccac tgggaatact gagcccatct ctccctacag 1200  
ctctctttac accagcattt ttttcacaga caccatcat actgactcca agcccttgc 1260  
tctccagtat ccacttctgg agtactctca gtctgttgc tcccctaagt ccagccagac 1320  
tgcaaggtgc taacacactt ttcagtttc ctctgtact gaacagtcac gggccattca 1380  
ctctgtctgg gctggatgga ccttccaccc ctggcccatt tccccagac ctacagaaga 1440  
25 cataacctat gcacttggg aatgagagaa ccgaggaaac aagaaacaga cattcaacat 1500  
gattgcattt gaagtgaaga acatctttt aggattctct ttgaatagga ctcaagtgg 1620  
tttgccattc ccattgaaa acatctttt aggattctct ttgaatagga ctcaagtgg 1680  
actatatgta taaaaatgcc ttaattggag tctaaactcc acctccctct gtcttttct 1740  
30 tttcttttc tttcttctt tcttcttct tctctctta aaaatatatt gagctttgtg 1800  
ctgaagaagt ttttgggtgg ctttagtgac tgtgtttgc aaaagcaatt aagaacaaag 1860  
ttactccttc tggctattgg gaccctttgg ccaggaaaaa ttatgcttag aatctattat 1920  
ttaaagaagt atttgtgaaa tgaaaaaaa aaaaaaaa aaaaaaaa 1933  
aaaaaaaaaaa aaa

35 <210> 21  
<211> 2215  
<212> DNA  
<213> Homo sapiens

40 <400> 21  
actcagtaga ccgccactgg ctgtgcacgt tatgggggtt ccacctaggg ctccgacctga 60  
ggcttgaac actccgtttt ccccccagtc acaggggagc tcttgcctcc cgcagctggg 120  
tcgcggtgtc tctcaaaagt cccctctac aggggcttcg tgaggcccg gcccacaggg 180  
45 cgctcggtcc cggaagtga gtctccaga ggggcccga gtggcagtg agggaggga 240  
gatggcggag gtgggggaga taatcgagg ctgccgcta cccgtgtgc ggcggaacca 300  
ggacaacgaa gatgagtgac ccctggcga gatcctgagc gtgaaggaca tcagtggccg 360  
gaagcttttc tacgtccatt acattgactt caacaaacgt ctggatgaat gggtagcga 420  
tgagcggctg gacctaaaga agatccagtt ccccaagaaa gaggccaaga ccccactaa 480  
50 gaacggactt cctgggtccc gtcttgctc tccagagaga gagggtcccg cctcggcgca 540  
ggccagcggg aagaccttgc caatcccggg ccagatcaca ctccgcttca acctgcccaa 600  
ggagcgggag gccattcccg gtggcgagcc tgaccagccg ctctcctcca gctcctgct 660  
gcagcccaac caccgctcaa cgaaacggaa ggtggagggt gtttaccag caactccagt 720  
gcccagcgag acagccccgg cctcgggttt tccccagaat ggagccgccc gtaggcgagt 780  
55 ggcagcccag ccaggacgga agcgaaaatc gaattgtttt ggcactgatg aggactccca 840  
ggacagctct gatggaatac cgtcagcacc acgcatgact ggcagcctgg tgtctgatcg 900  
aagccacgac gacatcgtca cccggatgaa gaacattgag tgcattgagc tgggcccggca 960  
ccgctcaag ccgtgtgact tctccccgta cccacaggaa ctccaccat tgctgtcct 1020  
ctacctgtgc gaggttctgc tcaagtacgg ccgtagtctc aagtgtcttc agcgtcattt 1080  
60 gaccaagtgt gacctacgac atcctccagg caatgagatt taccgcaagg gcaccatctc 1140  
cttctttgag attgatggac gtaagaacaa gagttattcc cagaacctgt gtcttttggc 1200  
caagtgttcc cttgaccata agacactgta ctatgacaca gacccttccc tcttctacgt 1260  
catgacagag tatgactgta agggcttcca catcgtgggc tacttctcca aggagaaaga 1320  
atcaacggaa gactacaatg tggctgcat cctaaccctg cctccctacc agcgcgggg 1380

- 18 -

ggtctctggtg gaacacagcc agacatcctg atggggaggc agagccagga agctaagcca 1320  
 gggcccagct gcgtccaacc cagcccccca cctcagggtc ctgaccccag ctcatgccc 1380  
 catcaccgcc tctccctggc tccccagggt gcagggtggc gcaaggccc gcccccatca 1440  
 catgttccct tggcctcaga gctgcccctg ctctcccacc acagccaccc agaggcacc 1500  
 5 catgaagctt ttttctcgtt cactcccaaa ccaagtgtc caaagctcca gtcctaggag 1560  
 aacagtccct gggtcagcag ccaggaggcg gtccataaga atggggacag tgggctctgc 1620  
 cagggtgccc gcacctgtcc agaacaacat gttctgttcc tctctctcat gcatttccag 1680  
 ccttg 1685

<210> 24  
 <211> 6428  
 <212> DNA  
 <213> Homo sapiens

15 <400> 24  
 gctagtggaa gttactgccg cgccaccgag tccggaccgg agactttggg gcctaactag 60  
 tgaatggtag tgtctagaaa gggatgtgct cttcaagaga gaggtgcaa tgtccaaccg 120  
 gcctaataac aatccagggg ggtcactgcy acgttcacag aggaacactg ccggggccca 180  
 20 accacaagac gactcaatag gaggaagaag ctgcagttca tcatctgctg tgatagtctc 240  
 acaaccagag gatccagaca gagccaatac ttcagaaaga caaaaaacgg ggcaggtgcc 300  
 taagaaagac aattctcgag gagtgaagcg cagtgcctagt ccagactaca acaggacca 360  
 ttctcctagc tctgcaaaaa aacaaaagc acttcagcat actgaatctc cctcagaaac 420  
 aaataagcca catagtaagt caaagaagag acatttagac caggagcaac aactgaaatc 480  
 25 tgcacaatca ccatcaacaa gcaaggctca taccaggaag agtggggcca ctggcggttc 540  
 acggagtcag aaaagaaaaa ggacagagag ttcttgtgta aagagtggct ccgggtctga 600  
 atcaactggt gcagaagaga gatctgcgaa acctaccaag ctggcttcaa aatcagccac 660  
 ctcagccaaa gctgggtgta gcaccatcac tgattcttct tctgctgctc ctacttctc 720  
 ctgctcttct gctgtagcct cggcctcctc cactgtacca ccaggtgcca gagtgaaca 780  
 30 aggaaaagat cagaacaagg ccaggcgttc ccgttcagcg tccagtccca gccccagaag 840  
 aagtagcagg gaaaaggaac agagtaaaac tgggtggctct tcaaaatttg attgggtctg 900  
 tcgtttcagc cctaaagtta gccttcctaa acaaaaactg tctcttccag ggtcttctaa 960  
 gtcagagaca tcaaaaacctg gcacttctgg attacaggcc aaattagcaa gtttaagaaa 1020  
 atctacgaag aaacgcagtg agtctccacc tgctgagctc cccagtttga ggcggagcac 1080  
 35 acgcaaaaag accacgggct cctgtgctag taccagtcgg cgaggctctg gcctgggcaa 1140  
 aagaggagca gctgaagctc gtcgacagga gaaaatggca gacctgaaa gcaaccagga 1200  
 ggcagtaaat tcttcaagctg ctcgacagga tgaagctccc caaggagctg caggggctgt 1260  
 tggcatgacc acctctgggg agagtgaatc agatgattcc gagatgggac gtttgcaagc 1320  
 tttgttagag gcaaggggct tccccctca cctatttggc cctcttggc ctcggtgtc 1380  
 40 acagcttttc catagaacaa ttggaagtgg agctagtctt aaggcccagc agctactaca 1440  
 aggattgcaa gccagtgatg aaagtcaaca gcttcaggca gttattgaga tgtgtcagtt 1500  
 actggtcatg ggaaatgagg agacactggg aggggttccct gtcaagagtg ttgttccagc 1560  
 tttgattacg ttacttcaga tggagcaca ttttgatatt atgaaccatg cttgtcagac 1620  
 cttaacatac atgatggaag cacttctctg atcttctgct gttgtagtag atgctattcc 1680  
 45 tgtcttttta gaaaagctgc aagtatttca gtgtattgat gtggcagagc aggccttgac 1740  
 tgccctggag atgttgctac ggagacatag taaagccatt ctacaggcgg gtggtttggc 1800  
 agactgcttg ctgtacctag aattcttcag cataaatgcc caaagaaatg cattagcaat 1860  
 tgcagctaatt tgctgccaga gtatcacgcc agatgaattt cattttgtgg cagattcact 1920  
 cccattgcta acccaaaagg taacacatca ggataaaaag tcagtagaaa gcacttgcct 1980  
 50 ttgttttgca cgcctagtgg acaacttcca gcatgaggag aatttactcc agcaggttgc 2040  
 ttccaaagat ctgcttcaaa atgttcaaca gctgttggta gtgactccac ccattttaag 2100  
 ttctgggatg tttataatgg tggttcgcat gtttctctg atgtgttcca actgtccaac 2160  
 tttagctggt caacttatga aacaaaacat tgcagaaacg cttcactttc tctgtgtgg 2220  
 tgccctcaat ggaagtgtgc aggaacagat tgatcttgtt ccacgaagcc ctcaagagtt 2280  
 55 gtatgaactg acatctctga tttgtgaact tatgccatgt ttacaaaag agggcatttt 2340  
 tgcagttgat accatgttga agaagggaaa tgcacagaac acagatgggt cgatatggca 2400  
 gtggcgtgat gatcggggcc tctggcatcc atataacagg attgacagcc ggatcattga 2460  
 gcaaatcaat gaggacacgg gaacagcacg tgccattcag agaaaacctc acccgtagc 2520  
 caatagtaac actagtggat attcagagtc aaagaaggat gatgctcgag cacagcttat 2580  
 60 gaaagaggat ccggaactgg ctaagtcttt tattaagaca ttatttgggt ttctttatga 2640  
 agtgatatgt tcttcagcag gacctgcggg cagacataag tgcccttagag caattcttag 2700  
 gataatttat tttgcggatg ctgaacttct gaaggatgtt ctgaaaaatc atgctgttcc 2760  
 aagtcacatt gcttccatgc tgtcaagcca agacctgaag atagtagtgg gagcacttca 2820  
 gatggcagaa attttaatgc agaagttacc tgatattttt agtgtttact tcagaagaga 2880

- 20 -

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 25

```

5  gcacgaggcc agccttacgg gcccgaaacc tcgtgtgaag ggtgcagtac ctaagccgga 60
   gcggggtaga ggcgggccgg caccctcttc tgacctccag tgccgcccgc ctcaagatca 120
   gacatggccc agaacttgaa ggacttgccg ggacggctgc ccgcccggcc ccggggcatg 180
   ggcacggccc tgaagctgtt gctggggggc ggccgctggt cctacggtgt gcgcgaatct 240
   gtgttcaccc tggaaaggcg gcacagagcc atcttcttca atcggatcgg tggagtgcag 300
10 caggacacta tcctggccga gggccttcac ttcaggatcc cttgggtcca gtacccatt 360
   atctatgaca ttcggggccag acctcgaaaa atctcctccc ctacaggctc caaagaccta 420
   cagatggtga atatctccct gcgagtgttg tctcgacca atgctcagga gcttcctagc 480
   atgtaccagc gcctagggct ggactacgag gaacgagtgt tgccgtccat tgttaatgag 540
   gtgtcaaga gtgtggggc caagttcaat gcctcacagc tgatcaccca gcgggcccag 600
15 gtatccctgt tgatccgccg ggagctgaca gagagggccca aggacttcag cctcatcctg 660
   gatgatgtgg ccacacaga gctgagcttt agccgagagt acacagctgc tgtagaagcc 720
   aaacaagtgg ccagcagga ggcacagcgg gcccaattct tggtagaaaa agcaaaagcag 780
   gaacagcggc agaaaattgt gcaggccgag ggtgaggccg aggtgccaa gatgcttga 840
   gaagcactga gcaagaaccc tggctacatc aaacttcgca agattcgagc agcccagaat 900
20 atctccaaga cgatcgccac atcacagaat cgtatctatc tcacagctga caaccttgtg 960
   ctgaacctac aggatgaaag ttaccacagg ggaagtgaac gcctcatcaa gggtaagaaa 1020
   tgagcctagt caccaagaac tccaccccca gaggaagtgg atctgcttct ccagtttttg 1080
   aggagccagc cagggggtcca gcacagccct acccgcccc agtatcatgc gatggctccc 1140
   cacaccgttt ccctgaaccc ctcttggttt aagggaagact gaagactagc cccttttctg 1200
25 ggggaattact ttcctctctc ctgtgttaac tggggctgtt ggggacagtg cgtgatttct 1260
   cagtgatttc ctacagtgtt gttccctccc tcaaggctgg gaggagataa acaccaacct 1320
   aggaattctc aataaatttt tattacttaa cctgaaaaaa aaaaaaaaaa 1370

```

30 &lt;210&gt; 26

&lt;211&gt; 642

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

35 &lt;400&gt; 26

```

   ccgcgcgtca cgtgacccca gcgcctactt gggctgagga gccgcgcgt cccctcgccg 60
   agtcccctcg ccagattccc tccgtcgccg ccaagatgat gtgcggggcg ccctccgcca 120
   cgcagccggc caccgcccag acccagcaca tcgccgacca ggtgaggtcc cagcttgaag 180
   agaaagaaaa caagaagtcc cctgtgttta aggccgtgtc attcaagagc caggtggtcg 240
40 cggggacaaa ctacttcac aaggtgcacg tcggcgacga ggacttcgta cacctgcgag 300
   tgttccaatc tctccctcat gaaaacaagc ccttgacctt atctaactac cagaccaaca 360
   aagccaagca tgatgagctg acctatttct gatcctgact ttggacaagg cccttcagcc 420
   agaagactga caaagtcac ctccgtctac cagagcgtgc acttgtgac ctaaaataag 480
   ctcatctccc gctgtgccct tggggtggaa ggggcaggat tctgcagctg cttttgcatt 540
45 tctcttccca aatttcattg tgttgatttc ttctctccc aatagggtgat cttaattact 600
   ttcagaatat tttaaaata gatattttt taaatcctt ac 642

```